

50 Years Since the Marr, Ito, and Albus Models of the Cerebellum

Mitsuo Kawato,^{a,b,*†} Shogo Ohmae,^{c†} Huu Hoang^a and Terry Sanger^{d,e,*}

^a Brain Information Communication Research Group, Advanced Telecommunications Research Institutes International (ATR), Hikaridai 2-2-2, "Keihanna Science City", Kyoto 619-0288, Japan

^b Center for Advanced Intelligence Project (AIP), RIKEN, Nihonbashi Mitsui Building, 1-4-1 Nihonbashi, Chuo-ku, Tokyo 103-0027, Japan

^c Department of Neuroscience, Baylor College of Medicine, Houston, Texas, USA

^d Department of Electrical Engineering, University of California, Irvine, 4207 Engineering Hall, Irvine CA 92697-2625, USA

^e Children's Hospital of Orange County, 1201 W La Veta Ave, Orange, CA 92868, USA

Abstract—Fifty years have passed since David Marr, Masao Ito, and James Albus proposed seminal models of cerebellar functions. These models share the essential concept that parallel-fiber-Purkinje-cell synapses undergo plastic changes, guided by climbing-fiber activities during sensorimotor learning. However, they differ in several important respects, including holistic versus complementary roles of the cerebellum, pattern recognition versus control as computational objectives, potentiation versus depression of synaptic plasticity, teaching signals versus error signals transmitted by climbing-fibers, sparse expansion coding by granule cells, and cerebellar internal models. In this review, we evaluate different features of the three models based on recent computational and experimental studies. While acknowledging that the three models have greatly advanced our understanding of cerebellar control mechanisms in eye movements and classical conditioning, we propose a new direction for computational frameworks of the cerebellum, that is, hierarchical reinforcement learning with multiple internal models.

This article is part of a Special Issue entitled: In Memoriam: Masao Ito—A Visionary Neuroscientist with a Passion for the Cerebellum. © 2020 The Author(s). Published by Elsevier Ltd on behalf of IBRO. This is an open access article under the CC BY license (<http://creativecommons.org/licenses/by/4.0/>).

Key words: Masao Ito, cerebellum, Marr, Ito, Albus.

INTRODUCTION

Masao Ito (1928–2018) made unique and profound contributions to our understanding of cerebellar functions, not only through experimental work (inhibitory action of Purkinje cells: Ito and Yoshida, 1964; Ito et al., 1964; Obata et al., 1967; adaptation of vestibulo-ocular reflex: Ito et al., 1982a; long-term depression of parallel-fiber-Purkinje-cell synapses: Ito et al. 1982b; Ito and Kano, 1982; molecular mechanisms of synaptic plasticity: Ito, 2001), but also through theoretical studies (Ito, 1970, 1974, 1984, 2006, 2008, 2011). Fifty years ago, David

Marr (Marr (1969)), Masao Ito (Ito (1970)), and James Albus (Albus (1971)) proposed computational models of cerebellar learning functions, based on neural circuits of the cerebellum (Eccles et al., 1967). As Ito acknowledged (Ito, 1970, 1974, 1984), his model was deeply influenced by Marr's model. However, the three models differed in several important respects.

With recent advances in experimental and computational studies, we compare and evaluate differences between these three seminal models (See Tyrrell and Willshaw, 1992; Sanger et al., 2020 for related efforts). In particular, we extend detailed discussions of three important differences of the models: parallel-fiber-Purkinje-cell synaptic plasticity, codon theory, and internal models. For oculomotor control and classical conditioning, Ito's model was sufficiently robust to influence experimental and computational studies for the past fifty years. While we acknowledge that the three models were among the most successful and influential computational models in neuroscience, they were insufficient as a computational framework for understanding several cerebellar functions, including whole body movements (Morton and Bastian 2007; Hoogland et al., 2015; Machado et al., 2015) or cognitive functions (Ito 2008; Strick et al., 2009; Lu

*Corresponding authors. Address: Brain Information Communication Research Group, Advanced Telecommunications Research Institute International (ATR), Hikaridai 2-2-2, "Keihanna Science City", Kyoto 619-0288, Japan.

E-mail addresses: kawato@atr.jp (M. Kawato), tsanger@uci.edu (T. Sanger).

[†] These two authors contributed equally to this work.

Abbreviations: AMPARs, α -amino-3-hydroxy-5-methyl-4-isoxazolepropionic acid receptors; IP₃, inositol 1,4,5-triphosphate; LTD, long-term depression; LTP, long-term potentiation; MAPK, mitogen-activated protein kinase; mGluR1, metabotropic glutamate receptor type 1; MOSAIC, Modular Selection And Identification for Control; STDP, spike-timing-dependent plasticity; IICR, IP₃-induced Ca₂₊ release; NMDAR, N-methyl D-aspartate receptor.

et al., 2012; Schmammann et al., 2019). In conclusion, we propose a new computational framework of hierarchical reinforcement learning with multiple internal models for phylogenetically newer parts of the cerebellum and newer functions.

COMPARISONS OF THE MODELS OF MARR, ITO, AND ALBUS

In this section we compare several characteristics of the three models proposed by Marr, Ito, and Albus (see Table 1). First, the three models all propose that modification of parallel-fiber-to-Purkinje-cell synapses guided by climbing-fiber inputs is central to sensorimotor learning in the cerebellum (Table 1). Ito acknowledged that his model followed Marr's model regarding this fundamental principle (Ito, 1970, 1974, 1984).

Neural circuits of the cerebellar cortex are much more uniform, regardless of different regions and zones (Sillitoe and Joyner 2007; Zhou et al., 2014; Tsutsumi et al., 2015, 2019), than those of the cerebral cortex. Based on this hardware uniformity, researchers, and especially theorists, are tempted to assume a uniform computational theory for the cerebellum (Apps and Garwicz, 2005; Apps and Hawkes, 2009). This assumption is not well substantiated for the following reasons. One of the outstanding contributions of David Marr (Marr (1982)) was his proposal of three levels of brain research: computational theory, representation and algorithm, and hardware implementation. If we accept that these three levels are somewhat independent, then hardware uniformity does not imply unified computational theory or uniform algorithms used in the cerebellum. Furthermore, different regions of the cerebellum receive mossy-fiber and climbing-fiber inputs from many different areas of the brain and the rest of the nervous system. Thus, input representations cannot be uniform either. Different regions of the cerebellum are involved in adaptation of oculomotor control, learning in conditioning, whole body movement, and cognitive functions. Correspondingly, mossy and climbing-fiber inputs range from direct inputs from sensory organs to inputs originating from higher association cortices of the cerebrum. Deep cerebellar nucleus outputs range from premotor neurons to higher cerebral association cortices via the thalamus. It would be very surprising if a single computational theory and a single representation and algorithm could explain all these diverse cerebellar functions, inputs, and outputs. Still, if we interpret the common proposal by Marr, Ito, and Albus as a computational algorithm used in the local circuits around Purkinje cells, some common algorithms and representations might be omnipresent in the cerebellum. We will explore mainly this possibility in this review.

Holistic versus modulating control

Marr and Albus proposed their models in a framework in which the cerebellum controls movements entirely by itself (holistic controlling role). Their models did not consider computational roles played by other brain regions or other parts of the nervous system, except for

climbing-fiber inputs. In contrast, Ito conceptualized the cerebellum in a network comprising other brain regions or other parts of the nervous system (complementary, modulating role, Table 1). For oculomotor control, conditioning, and higher cognitive functions, the cerebellum works as a side path in parallel to a main path formed by other brain regions or other parts of the nervous system.

Pattern recognition versus regression

The major objectives of most machine learning algorithms are either classification or regression. In the simplest classification, an algorithm determines whether a given sample belongs to a given class. In a regression, an algorithm approximates a target function using output computed from multiple inputs. In many cases of regression, the target function is smooth and nonlinear. Deep neural networks can classify millions of images into thousands of object categories, in what constitutes a pattern recognition problem. Three-layer neural networks can also approximate nonlinear inverse dynamics of multi-link robotic manipulators (Miyamoto et al., 1988), a regression problem that can be used for movement control.

Marr and Albus viewed pattern recognition tasks as objectives of cerebellar learning. In contrast, Ito targeted control tasks as objectives of cerebellar learning, thus treating them as regression problems. Adaptation of oculomotor control is best treated as a learning control and regression problem. Learning in a visually guided arm-reaching task likewise can best be treated as learning control and regression problems. In learning of conditioned responses, both pattern recognition views and control views are still advocated. Khilkevich et al., (2018) reported that over-trained rabbits with eyeblink conditioning either generate conditioned eyelid responses with a stereotyped amplitude, or else fail completely to generate eyelid responses, in an all-or-nothing manner. Since this pattern of response was maintained even when noise was added to mossy-fiber input, they regard this as experimental evidence that the cerebellum is able to recognize patterns whether mossy-fiber input is a conditioned stimulus or not. Thus, conceptually, the cerebellum is often interpreted as solving the pattern recognition problem of whether mossy-fiber inputs are conditioned stimuli. However, others still view the cerebellar role in conditioned learning as learning control, including amplitudes of conditioned responses, just as the terminology changed from “eyeblink conditioning” to “eyelid conditioning” (Medina et al., 2001, 2002). In summary, most sensorimotor learning problems are better viewed as control and regression problems, rather than pattern recognition and classification problems, because many experimental data demonstrate kinematic and kinetic representations of movements in the cerebellum. We must await future studies to see how much pattern recognition and how many regression characteristics are involved in the newest functions of cerebellar higher cognitive functions.

Table 1. Comparisons of the models of Marr, Ito and Albus. A check mark indicates an included concept, – denotes not mentioned, and a question mark means that the model was ambiguous on that point

	Marr	Ito	Albus
(1) Modification of parallel-fiber synapses on Purkinje cells conditioned by climbing-fiber inputs is central to learning	✓	✓	✓
(2) Holistic or Complementary role of cerebellum in motor control?	Holistic	Complementary	Holistic
(3) Pattern recognition versus Control	Pattern recognition	Control	Pattern recognition
(4) Synaptic plasticity of parallel-fiber-Purkinje-cell synapses when climbing fiber is activated	LTP	LTD	Anti-Hebbian
(5) Does climbing fiber input represent teaching signal, error signal or unconditioned stimulus?	Teaching signal	Error signal	Unconditioned stimulus
(6) Sparse expansion coding by granule cells or “Codon” theory	✓	—	✓
(7) Internal Models	—	✓	?

SYNAPTIC PLASTICITY IN ACHIEVING SUPERVISED LEARNING

Sites of synaptic plasticity for cerebellar learning

The three models assumed markedly different synaptic plasticity rules at parallel-fiber-to-Purkinje-cell synapses (Table 1). Marr assumed potentiation of these synapses when climbing-fibers, parallel fibers, and Purkinje cells are activated. Marr also postulated that in the cerebellum, only these synapses exhibit plasticity. In the past three decades, many different types of synapse have proven to be plastic (Table 2) (Hansel et al., 2001; Zhang and Linden, 2006; Hirano, 2013; D’Angelo et al., 2016), and many of them are Hebbian or anti-Hebbian (Gao et al., 2012). Among these, in addition to plasticity of molecular layer inhibitory interneurons, plasticity of mossy-fiber-to-deep-cerebellar-nucleus synapses has attracted experimental and theoretical interest intended to reveal their contributions to motor learning and its mechanisms (Miles and Lisberger, 1981; Lisberger and Sejnowski, 1992; Attwell et al., 2002; Tabata et al., 2002). Recently, memory transfer from parallel-fiber-to-Purkinje-cell synapses to mossy-fiber-to-deep-cerebellar-nucleus synapses has been intensively studied (Shutoh et al., 2006; Anzai et al., 2010; Yamazaki et al., 2015; Lee et al., 2015).

The number of parallel-fiber-to-Purkinje-cell synapses is 50 times larger than the sum of all other plastic synapses (Table 2). The capacity of the learning system, e.g., the allowable number of non-spurious memories in a recurrent artificial neural network (Hopfield, 1982; Gardner and Derrida, 1988), and approximation capacity of multi-layer perceptrons (Cybenko, 1989; Funahashi, 1989), increases as the number of parameters (Cover, 1965), or the number of synapses in biological networks, increases. Thus, learning capacity of the cerebellum largely depends on parallel-fiber-to-Purkinje-cell synapses, and it is reasonable that the three models emphasized these synapses so heavily. A huge learning system with trillions of learning parameters possesses immense learning capacity, but necessitates a huge learning sample for training, in order to guarantee good generalization in circumstances other than those in which the original learning occurred (Watanabe, 2009; Zhang et al., 2017; Suzuki, 2018; Schmidt-Hieber, 2017;

Table 2. Numbers of plastic synapses in the human cerebellum. The cell number column indicates the number of postsynaptic neurons. These numbers are based on Ito (1984), and Fukutani et al. (1992) and Andersen et al., (2004) for deep cerebellar nucleus

	Cell number	Synapse number
Parallel fiber – Purkinje	15 million	10,000 billion
Mossy fiber – Granule cells	50 billion	200 billion
Mossy fiber – Golgi cells	5 million	5 billion
Mossy fiber – deep cerebellar nucleus	500 thousands	5 billion
Purkinje cell – deep cerebellar nucleus	500 thousands	5 billion

Amari, 2020). A smaller learning system with a small number of synapses has a small learning capacity, but is better at generalization when provided with even a small number of learning samples. Transfer learning from parallel-fiber-Purkinje-cell synapses to mossy-fiber-deep-cerebellar-nucleus synapses may benefit from these respective advantages and disadvantages of huge and small learning systems. For example, long-term memory should possess more stringent generalization capability compared with short-term memory, because the former faces larger numbers of new different situations than the latter. Consequently, a small capacity system is more suited for long-term memory in situations in which only a fixed number of training samples is given (see Sanger et al., 2020 for related discussions).

LTD-LTP in parallel-fiber-Purkinje-cell synapses as supervised learning machinery

Ito proposed that climbing-fiber inputs represent error signals and that parallel-fiber-to-Purkinje-cell synapses undergo long-term depression (LTD) when both parallel-fiber and climbing-fiber inputs are activated (Table 1). Climbing-fiber inputs represent error signals originating from retinal slips in oculomotor control adaptation, including vestibulo-ocular reflexes, optokinetic responses, and ocular following responses (Simpson and Alley, 1974; Simpson et al., 1996; Kawano et al., 1996a, b; Kobayashi et al., 1998; Kawano, 1999). Especially for ocular following responses, parallel-fiber inputs represent

large visual-field motion in visual coordinates, while climbing-fiber inputs represent motor errors derived from retinal slips in motor coordinates (Kawano et al., 1996a,b; Kawato, 1999). This is a very rare example of an experimental paradigm for which sensory and motor coordinates are clearly separated. In visually guided arm reaching, climbing-fiber inputs represent errors (Kitazawa et al., 1998). Albus assumed that climbing-fiber inputs represent unconditioned stimuli, and that parallel-fiber-to-Purkinje-cell synapses obey the anti-Hebbian rule: synaptic efficacy decreases when parallel fibers, Purkinje cells, and climbing-fibers are all excited. In classical conditioning, climbing-fiber inputs can be interpreted as either unconditioned stimuli or unconditioned responses. However, this interpretation cannot be applied to broader classes of sensorimotor learning, and unconditioned stimuli or unconditioned responses can be equally well interpreted as sensory errors or motor errors, which can be reduced by learned conditioned responses.

Ito and others experimentally demonstrated LTD (Ito et al., 1982a,b; Ito and Kano, 1982; Hirano, 1990; Linden and Connor, 1995; Ito, 2001). Later, long-term potentiation (LTP) was discovered when parallel-fibers were activated in the absence of climbing-fiber inputs (Hirano, 1990, 2013; Sakurai, 1987). Accordingly, parallel-fiber-to-Purkinje-cell synapses possess bidirectional synaptic plasticity LTD-LTP controlled by the presence and absence of climbing-fiber inputs, respectively. Earlier there was some controversy as to whether LTD-LTP is essential for several types of motor learning in the cerebellum, but there is now a consensus that it is important and necessary (Welsh et al., 2005; Schonewille et al., 2011; Anzai and Nagao, 2014; Ito et al., 2014; Yamaguchi et al., 2016; Inoshita and Hirano, 2018; Kakegawa et al., 2018). If LTD-LTP is granted, it is contradictory to assign a teaching signal role to climbing-fiber inputs. Purkinje-cell simple-spike outputs would learn a sign-inverted waveform of the “teaching signal” that contradicts the “teaching signal.”

In the simplest interpretation, bidirectional LTD-LTP implements the Widrow-Hoff rule for ADALINE (Widrow et al., 1976; Fujita, 1982; Kawato et al., 1987; Kawato and Gomi, 1992) to minimize a squared error of regression in the steepest descent direction. The following is a mathematical demonstration that LTD-LTP can implement a simple supervised learning rule if the climbing-fiber firing modulation from its spontaneous level represents an error signal. Let us assume that Purkinje-cell simple-spike output y is a linear weighted summation of parallel-fiber input x_i by the parallel-fiber synaptic weight w_i ,

$$y = \sum_i w_i x_i$$

Purkinje-cell simple-spike output y learns to approximate a time-varying target value, \hat{y} , after sign inversion due to inhibitory action by Purkinje cells.

$$-y \rightarrow \hat{y}$$

If learning occurs in the steepest-descent direction of the squared error $E = (\hat{y} - (-y))^2 = (\hat{y} + y)^2$, the steepest-descent-direction of change in w_i is as follows,

$$-\frac{dE}{dw_i} = -\frac{d(\hat{y} + y)^2}{dw_i} = -2(\hat{y} + y)x_i$$

Let CF denote an instantaneous firing frequency of climbing-fiber input. \overline{CF} denotes its spontaneous firing level. The basic assumption implied in Ito (1970) was that if the difference $CF - \overline{CF}$ encodes the error $\hat{y} + y$, then LTD and LTP realize steepest-descent learning with a positive learning coefficient ε , as follows.

$$\delta w_i = -2\varepsilon(\hat{y} + y)x_i = -2\varepsilon(CF - \overline{CF})x_i$$

This equation is in agreement with LTD (negative δw_i , and w_i decreases) when $CF - \overline{CF}$ is positive (when climbing-fiber input is activated), and LTP (positive δw_i , and w_i increases) when $CF - \overline{CF}$ is negative (when climbing-fiber input is silent). Considering that the baseline or spontaneous climbing-fiber firing rate is low, around 1 Hz, the negative modulation below spontaneous firing rate often lead to a fall to zero. If $CF - \overline{CF}$ can be negative then δw_i is positive, when CF is smaller than \overline{CF} . In other words, LTP is triggered when climbing-fiber input is silent, which is less active than spontaneous firing. This means that the negative error signal would be an inhibition of spontaneous climbing-fiber inputs (Ohmae and Medina, 2015). This is the basic assumption of the unified LTP-LTD model (Kawato and Gomi, 1992).

Disinhibition of inferior olive nucleus neurons by Purkinje cells, via inhibition of the deep cerebellar nucleus, can implement the second term y of the error computation $\hat{y} + y$, at least partially (Nicholson and Freeman, 2003; Rasmussen et al., 2008; Chaumont et al., 2013; Ohmae and Medina, 2015). In order for this interpretation of LTD-LTP as the error minimizing synaptic-plasticity algorithm, the neural circuit that is downstream to deep cerebellar nucleus should have a proper sign and an aligned coordinate frame so that a change in $-y$ reduces the error $\hat{y} - (-y)$. These sign and coordinate requirements are fundamental to Ito's model, and were experimentally demonstrated at the detailed neural circuit level for oculomotor adaptation, including vestibulo-ocular reflex, optokinetic responses, and ocular following responses (Simpson and Alley, 1974; Ito, 1974, 2006; Gomi and Kawato, 1992; Kawano, 1999; Kawato, 1999; Yamamoto et al., 2002). For a broader class of sensorimotor learning by the cerebellum, examination of these theoretical requirements at the neural circuit level is still technically demanding, mainly because we do not fully understand the essential low dimensions at which learning occurs. We will discuss these error and dimension issues in the Section entitled, “Toward a new computational theory of the cerebellum.”

Coincidence detection between parallel-fiber and climbing-fiber inputs and a temporal window of synaptic plasticity

Doya (1999) postulated that the three learning algorithms proposed in computational neuroscience, unsupervised statistical learning, reinforcement learning, and supervised learning, are the main functions of the cerebral cor-

tex, the basal ganglia, and the cerebellum, respectively (see also [Raymond and Medina, \(2018\)](#) for cerebellar supervised learning). In the Section entitled, “Toward a new computational theory of the cerebellum”, we argue against this classification and propose that all three brain regions are for reinforcement learning. However, here we agree with [Doya \(1999\)](#), in the sense that a local neural circuit around Purkinje cells is most suited for supervised learning, based on a biophysical model of LTD for parallel-fiber-to-Purkinje-cell synapses ([Fig. 1A, B](#)). Large calcium increases in dendritic spines induce long-term decreases of synaptic efficacy, LTD, in the cerebellum ([Fig. 1C](#)), while it induces long-term increases of synaptic efficacy, LTP, in the cerebral cortex ([Fig. 1D](#)). In contrast, small calcium increases induce LTD in the cerebral cortex, while by themselves, they do not induce LTP in the cerebellum ([Tanaka et al., 2007](#)). Consequently, other factors such as nitric oxide are necessary for LTP induction in the cerebellum ([Ogasawara et al., 2007](#)).

[Fig. 1A](#) depicts schematically the early phase of LTD of parallel-fiber-Purkinje-cell synapses up to a large calcium increase ([Kuroda et al., 2001; Ito, 2002; Kotaleski et al., 2002; Doi et al., 2005](#)). Parallel-fiber inputs depolarize dendritic spines via α -amino-3-hydroxy-5-methyl-4-isoxazolepropionic acid receptors (AMPA). In parallel to this action, glutamate released from parallel fibers binds to metabotropic glutamate receptor type 1 (mGluR1) inducing a slow increase of inositol 1,4,5-triphosphate (IP₃) with 100 millisecond-order time to peak, via G-proteins and phospholipase C β . On the other hand, climbing-fiber inputs, which lag about 100 ms behind parallel-fiber inputs, induce large depolarizations in dendrites via multiple strong excitatory synapses, thereby opening voltage-dependent calcium channels on the spine and inducing calcium influx ([Ito, 2002; Doi et al., 2005](#)) ([Fig. 1A](#)). Because the latter electrical event is much faster than the former, IP₃ and Ca²⁺ concentrations increase simultaneously in the spine. This triggers a regenerative Ca²⁺ increase by IP₃-induced Ca²⁺ release (IICR) through IP₃-bound IP₃ receptors (IP₃Rs). IP₃Rs are IP₃-gated Ca²⁺ channels on the endoplasmic reticulum (ER), which is the intracellular Ca²⁺ store. IICR results in a supralinear Ca²⁺ surge with several micro-molar peaks ([Wang et al., 2000; Doi et al., 2005](#)) ([Fig. 1B](#)).

The Ca²⁺ surge induces subsequent reactions including activation of a mitogen-activated protein kinase (MAPK)-positive feedback loop, which phosphorylates and then internalizes AMPARs, and induces and consolidates LTD ([Kuroda et al., 2001](#)). [Tanaka et al., \(2007\)](#) demonstrated that a MAPK-positive feedback loop operates as a leaky integrator of Ca²⁺ concentration to determine the threshold of LTD, thus determining the temporal window of LTD. IP₃Rs and IP₃-dependent IICR act as coincidence detectors of the preceding parallel-fiber and delayed climbing-fiber inputs, as detailed in the following. Because the opening probability of IP₃Rs increases with IP₃ concentration, the Ca²⁺ threshold for IICR Ca²⁺ surge decreases within a few hundreds of milliseconds (blue curve in [Fig. 1B](#)) after multiple parallel-fiber inputs as IP₃ concentration

slowly increases via the mGluRs pathway, typically achieving its minimum after 100 milliseconds ([Fig. 1B](#)). Ca²⁺ increase by climbing-fiber inputs around this minimum-threshold time is sufficiently large to cross the threshold (thick black curve in [Fig. 1B](#)) so that IICR Ca²⁺ surge is triggered and drives LTD as explained above. If climbing-fiber inputs are either too early (gray curve in [Fig. 1B](#)) or too late (dotted curve in [Fig. 1B](#)) with respect to this minimum-threshold time, then the Ca²⁺ increase is below the threshold of the IICR Ca²⁺ surge and fails to induce LTD ([Fig. 1B](#)). In summary, the temporal window of LTD is determined by a slow increase in IP₃ concentration and the resultant dynamic threshold change determined by IP₃Rs kinetics, and is on the order of 100 milliseconds. Simplistically speaking, the molecular cascade consisting of glutamate, mGluR1, G-proteins, phospholipase C β , IP₃, IICR of IP₃Rs generates the delay such that the climbing fiber signal coming later can fall in the temporal window of LTD.

In addition, the intensity of climbing-fiber input can matter. Complex spikes measured in extracellular recording are not homogeneous and consist of 1–6 consecutive spikes of climbing-fibers (called spikelets; [Armstrong and Rawson, 1979; Maruta et al., 2007; Mathy et al., 2009](#)). The number of spikelets is essential to induce motor learning ([Rasmussen et al., 2013; Yang and Lisberger, 2014](#)). As the number of spikelets increases, it may reach the threshold of Ca²⁺ spikes, which could not be reached by shorter spikelets, and may extend the temporal window of Ca²⁺ surge and LTD. On the other hand, in saccadic adaptation, the timing of climbing-fiber input on the order of 100 milliseconds is suggested to be more important than the number of spikelets ([Herzfeld et al., 2018](#)). The ability of spikelet to extend the temporal window may vary, depending on the context and/or the area in the cerebellum.

Contrasting synaptic plasticity mechanisms between Purkinje cells and pyramidal neurons

[Fig. 1C, D](#) contrast different synaptic plasticity mechanisms of Purkinje cells of the cerebellum and pyramidal neurons of the cerebral cortex, while illustrating different morphological, physiological, and molecular mechanisms, with emphasis on coincidence detection. The large Ca²⁺ surge in Purkinje cells is induced by IICR from IP₃Rs and is mainly triggered by calcium influx caused by climbing-fiber inputs. Note that action potentials in Purkinje cells do not backpropagate because of excessive electrical load by extensive dendrite branching ([Vetter et al., 2001](#)) and the low density of sodium channels on dendrites ([Llinas and Sugimori, 1980; Stuart and Häusser, 1994](#)). Consequently, whether Purkinje cells are excited is irrelevant to LTD, while the large depolarization and subsequent Ca²⁺ increase induced by climbing-fiber inputs are decisive for LTD occurrence. Thus, supervised learning guided by error signals is suggested for the cerebellum. In contrast, in cerebral pyramidal neurons ([Fig. 1D](#)), NMDA receptors (*N*-methyl *D*-aspartate receptor, NMDAR) are coincidence detectors of glutamate released from presynaptic terminals and the backpropagating




Fig. 1. Different synaptic plasticity mechanisms of Purkinje versus pyramidal cells. **(A)** Schematic diagram of an LTD biophysical model (Doi et al., 2005). **(B)** Simulation reproduces the temporal window of Ca²⁺ surge that triggers LTD. **(C)** An LTD biophysical model overlaid on morphology, key molecules, and physiology of Purkinje cells. **(D)** Synaptic plasticity and coincidence detection mechanisms in pyramidal neurons in the cerebral cortex and medium ganglion cells in electrical lobes (ELL) in comparison with Purkinje cells in the same format as shown in **(C)**. *Figure **(A)** and **(B)** were published in *J Neurosci* 25, Doi T, Kuroda S, Michikawa T, Kawato M, Inositol 1,4,5-trisphosphate-dependent Ca²⁺ threshold dynamics detect spike timing in cerebellar Purkinje cells, 950–961, Copyright The journal of Neuroscience (2005). *Figure **(C)** and **(D)** were published in *Curr Opin Neurobiol* 21(5), Kawato M, Kuroda S, Schweighofer N, Cerebellar supervised learning revisited: biophysical modeling and degrees-of-freedom control, 791–800, Copyright Elsevier (2011).

action potential from the axon initial segment (Linden, 1999; Caporale and Dan, 2008). Glutamate, released from presynaptic terminals, binds to NMDARs, and backpropagating action potentials increase the postsynaptic voltage and consequently release a Mg²⁺-block of glutamate-bound NMDARs, resulting in full activation of NMDARs (Linden, 1999; Caporale and Dan, 2008; Urakubo et al., 2009). This leads to a large Ca²⁺ influx via NMDARs and induces subsequent reactions to consolidate LTP. Because the release of NMDARs from the Mg²⁺-block by backpropagating action potentials is the decisive event that leads to large calcium influx, Hebbian and unsupervised statistical learning are suggested for the cerebrum (Fig. 1D). For parallel-fiber-Purkinje-cell

synapses of adult cerebellum, there is no evidence that NMDARs in Purkinje cells participate significantly in LTD (Perkel et al., 1990; Renzi et al., 2007; Kono et al., 2019). In striatal medium spiny neurons, while Ca²⁺ influx depends on NMDAR activation by backpropagating action potentials, as in the cerebellum, synaptic plasticity also depends on activation of dopamine receptors (Wickens et al., 1996; Shen et al., 2008). In D1 receptor-expressing neurons, activation of the positive feedback loop, composed of PKA, PP2A and DARPP-32, serves as the coincidence detector of Ca²⁺ influx and dopamine input (Lindskog et al., 2006; Fernandez et al., 2006; Nakano et al., 2010); thus, the reinforcement learning rule is supported. Because D1 receptors and DARPP-32 are

expressed in prefrontal cortex, but at lower levels, the positive feedback loop probably does not possess bistability between the basal state and the excited state; thus, it cannot implement reinforcement learning rules, as in the basal ganglia. This is because any reinforcement learning rule necessitates memory functions provided by some kinds of bistability. In summary, Purkinje cell LTD is unique, supervised, and not Hebbian, while all those in the cerebral cortex, hippocampus, and basal ganglia are basically Hebbian, meaning that action potentials of the post-synaptic neurons are essential for synaptic plasticity.

According to the different mechanisms of coincidence detection in synaptic plasticity for the cerebellar cortex and the cerebral cortex, temporal windows of synaptic plasticity are also qualitatively and quantitatively different. The synaptic eligibility trace is a record of a synapse's recent activities to mark it as being eligible to change and to become distinct from other synapses (Barto et al., 1983). The synaptic eligibility trace for Purkinje cells is slowly rising IP_3 (Okubo et al., 2004; Doi et al., 2005); thus, the temporal window of LTD with respect to parallel-fiber input is ~ 100 ms in the cerebellum. In contrast, NMDAR dynamics are the synaptic eligibility trace for pyramidal neurons, dynamics that determine a temporal window of ~ 10 ms (Markram et al., 1997). Furthermore, when synaptic input precedes postsynaptic firing, LTP is induced, whereas if the synaptic input lags behind the firing, LTD is induced, which is spike-timing-dependent plasticity (STDP) (Bi and Poo, 2001). While late phases of LTD of the cerebellum and LTP of the cerebrum seem to share common signal-transduction mechanisms, the early phases are distinctly different, as explained above, and partly explain differences in learning algorithms at a micro level, as in dendritic spines.

Meta-plasticity of the temporal window of Purkinje-cell LTD

In the framework of Ito's model, simple spikes of Purkinje cells elicited by a combination of parallel-fiber inputs and spontaneous activities represent motor commands, and complex spikes of Purkinje cells elicited by climbing-fiber inputs represent error signals caused by executed movements. Thus, complex spikes should be delayed with respect to simple spikes, because of feedback delays. This delay includes latency for conduction of command signals within a neural circuit downstream to the cerebellum, muscle stretches, movement execution, sensory organ activation, afferent signal conduction, and finally activation of inferior olive neurons. Yamamoto and colleagues (2002) assumed that this delay was compensated by the temporal window of LTD and conducted efficient learning simulations. That is, the temporal correspondence between the responsible motor command (parallel-fiber inputs and simple spikes) and the resulting motor error (climbing-fiber inputs and complex spikes) is guaranteed in LTD, if the motor-sensory time delay is matched to the temporal window of LTD (Ogasawara et al., 2008). The motor-sensory time delay should be different for different cerebellar regions controlling different movements. For example, the delay is expected to be small, around 30 ms, for ventral paraflocculus controlling


ocular-following responses. The delay is expected to be large and longer than 100 ms for a region in a hemisphere responsible for visually guided arm-reaching. The size (width and center) of the LTD temporal window for different parts of the cerebellum should be matched to the motor-sensory time delays, if the objective of the LTD temporal window is to cancel the delay. Suvrathan and colleagues (2016, 2018) found experimental data supporting this theoretical requirement. In the biophysical model of LTD (Doi et al., 2005), the optimal time delay between parallel-fiber inputs and climbing-fiber inputs for effective LTD, the time for IP_3 concentration to reach its maximum after parallel-fiber inputs arrive, is determined by a biochemical reaction delay in the mGluR pathway. This biochemical delay depends on concentrations of the molecules involved, three-dimensional arrangements of anchor proteins, diffusion constants of molecules, and geometrical characteristics of cytosols within dendritic spines. It would be exciting research to theoretically and experimentally investigate possible chemical mechanisms for meta-plasticity to change some of the above factors so that the biochemical delay is matched to the neural delay in a region- and movement-specific manner in the cerebellum.

CODON THEORY

Marr and Albus proposed "Codon Theory" for representations and computations by granule cells, while Ito did not address this subject (Table 5). Because Brindley (1964, 1969), a mentor of Marr, already discussed cerebellar-supervised learning, codon theory is one of the most innovative elements of Marr's model. The number of cerebellar granule cells is ~ 50 billion, exceeding the sum of all other neurons in the brain. Each granule cell possesses 4 to 5 synapses on small dendrites and receives synaptic inputs from mossy-fibers. The number of granule cells is 200 times the number of mossy-fibers, ~ 250 million (Fig. 2B). This corresponds to "expansion coding" in modern terminology. In modern computational terminology, codon theory also postulates "sparse coding" such that only a small number of granule cells are activated when specific combinations of their mossy-fiber inputs are activated. By expanding the dimension of input representation (expansion) and reduction of the number of simultaneously activated granule cells (sparseness), interference of associative memories for different contexts is minimized; thus, efficient learning can be expected.

Codon theory predicts nonlinear input–output transformation at granule cells, low firing rates of granule cells, activation of only a small portion of all granule cells, and multiple modalities of mossy-fiber inputs to individual granule cells, at least for certain regions of the cerebellum and for certain cerebellar functions. In the past two decades, large technical advances of intracellular and patch clamp recordings, as well as Ca^{2+} imaging from granule cells have been made. Contradictory data for the above four predictions has been obtained and some have even argued that codon theory has been disproven. Sanger et al., (2020)

A



B




Fig. 2. (A) Circuit diagram of the cerebellum with the dentate nucleus, emphasizing its parallel path characteristics to the cerebral cortex in sensorimotor control (taken from Sanger et al., 2020). In (A), IO: inferior olive nucleus, CPC: cortico-ponto-cerebellar pathway, CTC: cerebello-thalamo-cortical pathway. **(B)** Relative numbers of cells and synapses in mossy-fibers, granule cells, Purkinje cells, and dentate-nucleus circuits (taken from Sanger et al., 2020). In (B), the area of each gray rectangle is proportional to the approximate cell count for the corresponding cell type. The area of each blue inset illustrates the relative number of output synapses from that type of neurons. *Fig. 2 was published in Journal of Physiology, 598(5), Sanger TD, Yamashita O, Kawato M, Expansion coding and computation in the cerebellum: 50 years after the Marr-Albus codon theory, 913–928, Copyright John Wiley and Sons (2020).

recently reviewed this evidence and pointed out that seemingly contradictory data obtained under specific preparations and conditions cannot disprove the core of codon theory. Because of this recent review, here we only very briefly discuss the above four points: nonlinear transformation, low firing rates, sparse coding, and multiple input modality.

Under some experimental conditions, granule cells can fire at high rates when activated by only a single mossy-fiber, and the input–output firing-rate relationship is linear (Jörntell and Ekerot, 2006; Rancz et al., 2007; Arenz et al., 2008; van Beugen et al., 2013). Kawano and colleagues showed that instantaneous firing rates of both mossy-fibers and parallel fibers are related to eye movement dynamics and are similar during ocular following responses in ventral paraflocculus (Kawano and Shidara, 1993; Kawano et al., 1996a,b). Head rotation velocities were predicted from mossy-fiber firing frequencies to granule cells, suggesting that instantaneous firing rates encode information in granule cells (Arenz et al., 2008). Recent Ca^{2+} imaging studies indicated that a large proportion of granule cells are excited simultaneously, which appears to contradict sparse coding (Spanne and Jörntell, 2015; Badura and De Zeeuw, 2017; Giovannucci et al., 2017; Knogler et al., 2017; Wagner et al., 2017; Gilmer and Person, 2018). However, this point critically depends on temporal resolution, or on a minimal meaningful time bin of granule cell encoding. In some examples of eye movements, temporal resolution of simple-spike firing is around 1 millisecond (Shidara et al., 1993; Gomi et al., 1998; Payne et al., 2019). If a sampling frequency of Ca^{2+} imaging is 30 Hz and 50% of granule cells are co-activated in the imaging resolution, minimally only 1.5% of granule cells are simultaneously activated in a 1-millisecond time bin. For the vestibular cerebellum and regions that directly receive tactile inputs, under decerebrated and/or anesthetized conditions, granule cells receive synaptic inputs from a single modality (Jörntell and Ekerot, 2006; Bengtsson and Jörntell, 2009). However, Ishikawa et al., (2015) found that mossy-fiber inputs of somatosensory, auditory, and visual modalities from the cerebral cortex converge on individual granule cells (Chabrol et al., 2015; Shimuta et al., 2019). In this case, nonlinear input–output relationships were observed for combined stimuli. Anatomically, one mossy-fiber carrying proprioception from the external cuneate, and another mossy-fiber carrying other modalities from pontine nuclei converge to a single granule cell (Huang et al., 2013).

In the Section, Comparisons of models by Marr, Ito, and Albus, we discussed pattern recognition versus control functions of the cerebellum. An all-or-nothing (firing or not) representation is best suited for pattern recognition, and codon theory is based on these assumptions. Even for instantaneous firing-rate coding and control (regression), codon theory might be relevant to enhancement of approximation precision in regression, as follows. Neuronal origins of mossy-fibers, granule cells, and Purkinje cells form a three-layer feedforward neural network. Funahashi (1989) and Cybenko (1989) mathematically proved that such three-

layer neural networks can learn to approximate any arbitrary, smooth, nonlinear functions if the intermediate layer has a sufficiently large number of neurons, and synaptic weights are optimally chosen. As will be discussed in the Section, Internal models, function approximation capability is essential for learning acquisition of forward models or inverse models of nonlinear, controlled objects within the three-layer network. Because mossy-fiber-granule-cell synapses possess Hebbian plasticity (Schweighofer et al., 2001; Sgritta et al., 2017), backpropagation learning algorithms used for deep neural networks cannot be applied. Consequently, in order to increase function approximation capability of the three-layer network, a huge number of granule cells and a rich repertoire of nonlinear basis functions are essential there (Sanger et al., 2020). In this sense, the spirit of codon theory, which was proposed in pattern recognition domains, is still relevant even for cerebellar functions and regions related to control, regression, and instantaneous firing-rate coding. This is because the rich repertoire of nonlinear transformations, provided by a vast number of granule cells, is essential for approximation capabilities by the cerebellum, and it is a core assumption of codon theory.

Higher expansion ratios are beneficial for more elaborate cerebellar functions dealing with strong nonlinearity; thus, the relevance of codon theory is higher for larger expansion ratios. Recent structural magnetic resonance imaging studies on gray matter and input fibers of human cerebellum suggested that expansion ratios of granule cells from mossy fibers are different between the vermis and hemispheric regions. The gray-matter volume of the hemisphere was 11.4× of that of the vermis, while the cross-sectional area of the cerebrocerebellar peduncle was estimated at less than 2.8× the spinocerebellar tract (Keser et al., 2015). On the assumption that diameters of mossy-fibers of these two regions of the cerebellum do not differ, we estimate that the expansion ratio of the hemisphere is 4x larger than that of the vermis. Large expansion ratios might be more marked for phylogenetically newer parts of the cerebellum and higher cognitive functions. In contrast, seemingly contradictory data against codon theory were mostly obtained in older parts of the cerebellum under anesthetized or decerebrate preparations. We postulate that codon theory is more relevant to newer functions of the cerebellum.

INTERNAL MODELS

Cerebellar internal models

Internal models in neuroscience are the neural networks that simulate input–output relationships of some processes, such as controlled objects in movement execution. “Internal” implies that the neural network is within the brain or the cerebellum. “Model” implies that the neural network simulates a target dynamical process. When humans move their bodies quickly, the necessary computation is too difficult to be solved by simple feedback controllers alone, and internal models are necessary (Gomi and Kawato, 1996). Here, we

include both forward and inverse (see below) models as possibilities. There is always feedback control, but except very basic feedback control, model-predictive controllers or precomputed ballistic movements include some forms of internal models. When humans grow their bodies and manipulate different objects, their dynamics change drastically; thus, internal models cannot be genetically preprogrammed or fixed, and must be acquired through learning. Ito (1970) proposed that internal models are acquired by learning in the cerebellum (Table 1; Fig. 3A, B). Neither Marr nor Albus mention internal models around 1970, but Albus (1975) later proposed an artificial neural network model called CMAC (cerebellar model articulation controller), and one of its possible applications was to learn inverse dynamics models of robots.

Internal models are classified as either forward or inverse. Forward models possess the same input–output direction as controlled objects and simulate their dynamics. For example, controlled objects such as eyeballs or arms receive motor commands and generate movement trajectories, that can be represented as sensory feedback about the executed trajectories. Forward models receive copies of motor commands (corollary discharge or efference copy) and predict movement trajectories or sensory feedback (sensory consequence). In control engineering and robotics, forward models have been and still are just called “internal models.” Jordan and Rumelhart (1992) coined this term to discriminate them from inverse models, and this terminology was soon adopted by cognitive science and neuroscience. On the other hand, inverse models simulate inverted input–output relationships of controlled objects. In a sense, inverse models provide inverse functions of modeled dynamical systems. Inverse models of controlled objects can receive desired trajectories as inputs and can compute necessary motor commands to realize the desired trajectories. Because an inverse model possesses the inverse input–output relationship of a controlled object, a cascade of the inverse model and the controlled object becomes an identity map. This implies that the trajectory realized by this cascade is equal to the desired trajectory; thus, the inverse model provides an ideal feedforward controller. We note that forward models are well-posed problems (every action causes a unique result) whereas inverse models are not (the same result can occur following many actions). So, among many possibilities, the brain needs to select one for the inverse model to learn. Although the combination of an inverse model followed by a forward model is the identity, it is not unique.

Fig. 3C, D show our interpretations of Fig. 3A, B proposed by Ito, while utilizing modern terminology and computational backgrounds of inverse and forward models, respectively. In Fig. 3C, the 3-neuron vestibulo-ocular reflex arc and the flocculus form a parallel system and the combined system provides an inverse dynamics model of a controlled object, a downstream of the oculomotor system from the vestibular nucleus. Note that the 3-neuron arc is fixed and the flocculus is modifiable; thus, the combined system is modifiable and can learn the inverse dynamics model. The sensory

error between the desired trajectory (sign inverted head rotation) and the realized trajectory (eye rotation) is given as a retinal slips and it is transformed into motor coordinates by the control network of optokinetic responses, and used as the motor command error to guide learning of a part of the inverse dynamics model in the flocculus (Gomi and Kawato, 1992; Tabata et al., 2002). Fig. 3B can be also interpreted to mean that a newer part of the cerebellum provides an inverse dynamics model of a controlled object. In this interpretation, a sensorimotor cortex of the cerebrum provides a feedback motor command produced through long-loop sensory feedback through an external world, and this feedback motor command is utilized to train the inverse dynamics model in the cerebellum, that is, it functions as an ideal feedforward controller. In another interpretation based on the forward model (Fig. 3D), part of the cerebellar hemisphere provides a forward model of the controlled object. It computes a predicted trajectory while receiving the efference copy of the motor command. The difference of the realized trajectory and the predicted trajectory provides a sensory prediction error that can be used as an error signal to train the forward model. This forward model can be used as an essential element for internal feedback control, bypassing the long-loop through the external world, reducing feedback delays and increasing control performance. Fig. 3D illustrates this latter interpretation of Fig. 3B based on later computational models of internal feedback control with forward models (Kawato et al., 1987; Kawato and Gomi, 1992; Miall et al., 1993). A linear quadratic regulator (Bemporad et al., 2002; Li and Todorov, 2004) and an optimal feedback controller proposed by Todorov and Jordan (2002) can be regarded as more sophisticated feedback control algorithms, which also utilize forward models.

As discussed in the Section, Codon theory, the 3-layer feedforward neural circuit comprising neuronal origins of mossy-fibers, granule cells, and Purkinje cells, is expected to closely approximate arbitrarily complex nonlinear functions with a vast number of granule cells, if an appropriate error signal is provided and optimal synaptic weights are learned. Thus, the cerebellum basically possesses learning capability to acquire internal models if climbing-fiber inputs provide appropriate error signals. Kawato and colleagues (1987) proposed that forward and inverse models are acquired distinctively in different parts of the cerebellum. Learning of forward models is computationally straightforward, since the error signal can be computed as the difference between the realized trajectory (or sensory feedback) and the predicted trajectory (or predicted sensory feedback) (Fig. 3D) (see also discussion of computing $\hat{y} + y$ by Purkinje-cell disinhibition in the Section, Synaptic plasticity in achieving supervised learning). In contrast, learning of inverse models is fundamentally difficult, because we cannot assume the existence of a teaching signal \hat{y} , i.e., ideal motor commands in the brain. This is because if such an ideal motor command already exists, it can be used as the appropriate motor command in movement executions, and inverse models or learning them is unnecessary. We postulated a “feedback-error-learning”




Fig. 3. Figures from Ito (1970) and interpretations thereof, based on inverse and forward internal models. (A) and (B) are two figures from Ito (1970), and (C) and (D) are their interpretations, respectively. (A) Fig. 6(C) of Ito (1970) for feedforward control of the vestibulo-ocular reflex. FL, flocculus, V, sensory input from the vestibular organ, SV, rostral part of the vestibular nuclear complex, OM, oculomotor neuron, EB, eyeball, TL, teaching line for the learning in cerebellar cortex. (B) Fig. 7 (B) of Ito (1970) for the control system of voluntary movements. AC, cerebral association area, MC, cerebral motor area, SM, spinal motor system, SC cerebral sensory area, MA, motor activity, H, internalized feedback pathway through the external world, NC, neocerebellum in which SM, MA, H, SC, and AC are modeled in a miniaturized form. (C) Inverse dynamics model interpretation of (A) for adaptation of the vestibulo-ocular reflex. FL and V-SV-OM in (A) correspond to flocculus and 3 neuron arc in (C), respectively. Dashed rectangles in (A) and (C) correspond with each other. The flocculus and the 3-neuron arc function in parallel to acquire an inverse dynamics model of the controlled object. A feedback controller for optokinetic responses in the accessory optic system transforms a retinal slip into the motor error, which is used for adaptive changes of the inverse dynamics model in the flocculus. (D) Forward internal model interpretation of (B). The neocerebellum acquires a forward model of a controlled object, and uses it for internal feedback control. NC in (B) corresponds to forward model in (D). MC in (B) corresponds to feedback controller in (D). H in (B) corresponds to upper feedback loop in (D).

scheme in which the inverse dynamics model receives the error signal in a motor command coordinate from the output of the feedback controller, which performs more crude and poorer sensorimotor transformation than the inverse model can do, but still provides the necessary motor error (Kawato et al., 1987). The feedback controller receives a sensory error as its input, and the output from the feedback controller can be regarded as the motor error necessary for training the inverse model (Fig. 3C). Accordingly, climbing-fiber inputs for some regions were postulated to represent outputs of feedback controllers (Kawato and Gomi, 1992; Gomi and Kawato, 1992; Bhushan and Shadmehr, 1999; Tabata et al., 2002; Ramnani, 2006; Ito, 2008). Inoue and colleagues (2016, 2018) showed that motor commands of cerebellar motor cortex in non-human primates can drive motor adaptation and can be potential sources of error signals for the cerebellum, in accordance with the feedback-error-learning scheme.

Experimental support for forward and inverse models

Experimentally, it is not easy to address whether the cerebellum is predicting the sensory consequence of the movement (forward model) or computing the necessary feedforward motor command (inverse model). Fundamental mathematical difficulties in asking this question were demonstrated by Mehta and Schaal (2002). It has been reported that the activity of cerebellar Purkinje cells and cerebellar nuclei encode kinematics (position and/or velocity) of movement, suggesting forward models in the cerebellum (Laurens et al., 2013; Brooks and Cullen, 2013; Brooks et al., 2015; Herzfeld et al., 2015, 2018; Streng et al., 2018). However, since signals of position or velocity can be a part of the motor command, the possibility of signaling by inverse models cannot be excluded. Similarly, some experimental data demonstrated that Purkinje cell activities directly drive movements and sup-

port inverse models rather than forward models (Heiney et al., 2014; Bhanpuri et al., 2014; Hoogland et al., 2015; Lee et al., 2015; El-Shamayleh et al., 2017; Chabrol et al., 2019; Payne et al., 2019). However, it is also possible that stimulation interferes with normal prediction of sensory consequences of movement and induces artifactual movements by sensori-motor transformation in the downstream of the cerebellum. To solve the problem of discriminating forward from inverse models, carefully designed paradigms are required to dissociate movement consequences and motor commands. Pasalar et al., (2006) and Yamamoto et al., (2007) are representative examples of such studies in non-human primates, but surprisingly, they drew opposing conclusions. The former supported the forward model, while the latter supported the inverse model. This discrepancy can be explained by assuming that slow movement in the former study does not require feedforward control by an inverse model, whereas feedforward control was essential for much faster movements in the latter study. Although other research also supported cerebellar forward models (Wolpert et al., 1995; Blakemore et al., 2001; Kawato et al., 2003; Bastian, 2006; Tseng et al., 2007; Izawa et al., 2012; Sultan et al., 2012), experimental data for cerebellar inverse models are more straightforward in their interpretations. In particular, various studies on ocular following responses in the ventral parafovea (see the Section, Synaptic plasticity in achieving supervised learning) have demonstrated that Purkinje cell input and output correspond to sensory and motor coordinates, respectively, and that the inverse model is computed in the cerebellum (Shidara et al., 1993; Kawano et al., 1996a,b; Gomi et al., 1998; Kobayashi et al., 1998; Kawano, 1999; Kawato, 1999; Yamamoto et al., 2002; Takemura et al., 2017).

Electrosensory lobe and forward models

One of the important functions of forward models is central cancellation of sensory consequences of autogenous movement. The electrosensory lobe of weakly electric fish is one of many cerebellum-like structures, and illustrates beautifully this proposed function. Medium ganglion cells of the electrosensory lobe estimate electrosensory signals induced by external events by subtracting electrosensory signals caused by the fish's own electric organ discharge and movements (Bell and Russell, 1978; Han et al., 2000; Requarth and Sawtell, 2011; Kennedy et al., 2014; Sawtell, 2017). This can be regarded as a special case of estimating sensory inputs. Forward models can predict sensory inputs caused by an animal's own movement from the efference copy of motor commands. Because of this finding, some argue that the cerebellum should also cancel sensory inputs caused by autogenous movements utilizing forward models, i.e., central cancellation of sensory consequences of autogenous movements by forward models. However, this expectation is not fully supported when we consider different synaptic plasticity mechanisms between medium ganglion cells and Purkinje cells and different representations of climbing-fiber inputs to different types of neurons. In Fig. 1C, D, we con-

trasted different synaptic plasticity mechanisms of Purkinje cells and pyramidal neurons of the cerebrum. Synaptic plasticity rules of medium ganglion cells are anti-Hebbian and are more similar to those of pyramidal neurons than of Purkinje cells. We note that anti-Hebbian learning requires action potential generation in post-synaptic neurons for depression of synapses, which is different from Purkinje cell LTD. As already explained, action potentials do not backpropagate in Purkinje cells, and so far there is no evidence that post-synaptic firing in Purkinje cells contributes to the synaptic plasticity of LTD. Climbing-fiber inputs to medium ganglion cells represent total electrosensory signals, and parallel-fiber inputs include motor corollary discharge signals related to the electric organ discharge, and proprioceptive signals related to movements and position of the tail, trunk, and fins. When the medium ganglion cell, and parallel-fiber synapse carrying the corollary discharge, and climbing-fiber input are all simultaneously activated, according to anti-Hebbian learning, parallel-fiber synapses undergo LTD. Consequently, learning cancels the electrosensory signal that can be predicted by the corollary discharge from the firing output of the medium ganglion cell. As explained in the Section, Synaptic plasticity in achieving supervised learning, this anti-Hebbian learning is completely different from the supervised learning rule of Purkinje cells, and the same cancellation computation is impossible for Purkinje cells. If climbing-fiber inputs to Purkinje cells encode the error between the actual sensory input and that predicted by the corollary discharge, then climbing-fiber inputs should decrease to zero while Purkinje cells continue to fire, if there is no external sensory event after learning is completed. In contrast, climbing-fiber inputs to medium ganglion cells continue to fire while medium ganglion cells become silent when there is no external sensory event after learning is completed. Accordingly, both upstream and downstream neural circuits represent markedly different pieces of information, so they should be wired entirely differently for the cerebellum and the electrosensory lobe. From computational viewpoints, Purkinje cells can perform sensori-motor transformation, whereas medium ganglion cells simply subtract climbing-fiber inputs from parallel-fiber inputs within the same dimensions without coordinate transformation. We acknowledge that some parts of the cerebellum most probably acquire forward internal models for central cancellation of sensory consequences of autogenous movement (Blakemore et al., 1998, 2001; Wolpert et al., 1998; Kawato et al., 2003; Brooks and Cullen, 2013; Brooks et al., 2015), but experimental and computational evidence from the electrosensory lobe is not directly relevant to this function of the cerebellum.

Cerebellar internal models for cognition

Cerebellar internal models can simulate any dynamical processes other than controlled objects in sensorimotor tasks, but only if climbing-fibers represent appropriate error signals between the actual and predicted outputs from the dynamical processes, and possibly provide computational mechanisms for recent proposals of

cerebellar cognitive functions (Ito, 2008; Strick et al., 2009; Lu et al., 2012; Honda et al., 2018; Schmahmann et al., 2019; Devereitt et al., 2019). Higher cognitive functions are realized in the brain, utilizing cerebral cortex, basal ganglia, and subcortical structures. If inputs and outputs of these computational modules are provided to newer parts of the cerebellum as parallel-fiber and climbing-fiber inputs, different parts of the cerebellum can learn to acquire either forward or inverse models of these cognitive processes. Imamizu and Higuchi showed that internal models of many different tools are acquired in the human cerebellum in a modular manner, and that their representations largely overlap with those for language in the cerebellum, as well as in Broca's area of the cerebral cortex (Imamizu et al., 2000, 2003; Higuchi et al., 2007, 2009). If multiple internal models are acquired in a modular and switching mode suggested by mixture-of-experts model of artificial neural networks (Jacobs et al., 1991), they are advantageous to cope with very complicated tasks such as rapidly switching manipulation of many different objects (Gomi and Kawato, 1993). A basic form of Modular Selection And Identification for Control (MOSAIC) architecture consists of multiple paired forward and inverse models, and can be regarded as a more parallelized version of the mixture of experts architecture (Wolpert and Kawato, 1998). Each forward model tries to predict local dynamics and competes with other forward models for goodness of prediction, which is measured as a responsibility signal. The product of the error and the responsibility signals is given as the final error signal for training the forward model and the corresponding inverse model (Fig. 5B). A "divide and conquer" learning strategy is implemented in this parallel architecture. A prior probability of the responsibility signal can be learned by a responsibility predictor. The responsibility predictor can be interpreted as a parallelized version of the gating network of the mixture-of-experts architecture. MOSAIC can work either in supervised (Fig. 5B) or reinforcement learning (Fig. 5A) paradigms (Haruno et al., 2001; Doya et al., 2002). In the case of reinforcement-learning MOSAIC, an actor or a critique or both take the role of the inverse model. Because of the dualistic nature of forward and inverse models, MOSAIC can generate a sequence of complicated behaviors, and can also recognize patterns in symbolic representations of the sequence by observing continuous movement trajectories (Kawato et al., 2000; Wolpert et al., 2003; Samejima et al., 2006; Kawato and Samejima, 2007). Based on these experimental and theoretical explorations of possible computational roles of cerebellar internal models in cognitive functions, Ito (2008) proposed a cognitive framework of cerebellar internal models for thought processes. Fig. 4 shows the number of PMC papers published per year with the following three PubMed search conditions; {(internal model) AND cerebellum}, [(internal model) AND cerebellum] AND (motor control), [(internal model) AND cerebellum] AND cognition. Rapid increases in all three categories after 2000 indicate that cerebellar internal models for both motor control and cognition have begun to be widely accepted in the last two decades.

TOWARD A NEW COMPUTATIONAL THEORY OF THE CEREBELLUM

The three models proposed by Marr, Ito, and Albus were extremely successful in establishing a motor learning framework within which our understanding of the cerebellum was advanced for oculomotor control, classical conditioning, and visually guided arm-reaching. However, these models are not sufficient to extend our understanding of cerebellar functions to whole body motions or cognition, because of computational and experimental reasons that we will discuss in this section. Furthermore, based on analyses of difficulties with the three models, we propose a new computational framework of the cerebellum.

The curse of dimensionality

First, the three models do not provide insights regarding how to resolve "curse of dimensionality" associated with learning in systems with many degrees of freedom. Numbers of muscles and neurons used by animals in whole body movements are huge. Machine learning algorithms, including supervised and reinforcement learning, confront difficulties with such large systems. If a whole-body movement includes 1000 muscles, and each muscle can generate 100 different levels of force, a simple and plain reinforcement learning algorithm needs to visit and explore on the order of 100^{1000} states to find a good solution. For example, unsatisfactory results from the final competition of the DARPA robotic challenge demonstrated these difficulties even with systems having far fewer degrees of freedom than biological systems, that is, humanoid robots (Atkeson et al., 2018). Under some mathematical assumptions, a statistical learning theory estimates a generalization error for a test dataset not used for training, as $d/2n$, where d is the number of parameters in the classification algorithm and n is the number of training samples (Watanabe, 2009). Recent asymptotic theories motivated by the success of deep neural networks estimated that the number of necessary training samples for good generalization could be relaxed, compared with the above estimate (Zhang et al., 2017; Schmidt-Hieber, 2017; Suzuki, 2018; Amari, 2020), but tens of millions of training samples are still needed for tens of millions of synapses. Considerations of these theoretical constraints for cerebellar learning suggest an astronomical number of necessary training samples to achieve good generalization, with unacceptably long training periods, as follows. Let us assume that there exist 10,000 microzones in the human cerebellum, and that each microzone works as a single machine-learning entity. Then each microzone possesses 1 billion parallel-fiber-Purkinje-cell synapses (Table 2). To attain a 5% generalization error (that is, 95% correct), the above estimate requires 10 billion training samples. In extremely fast sampling, let us assume that a learner collects training data at 100 Hz (1 sample per 10 ms), 3600 sec per hour, 8 hours per day. Under these assumptions, 347 days are necessary to collect 10 billion samples. This extremely underestimated necessary training period is




Fig. 4. Numbers of PMC papers published each year, as shown by PubMed searches for the keywords “cerebellum and internal models”, and “motor control” or “cognition”. The orange line indicates [internal model AND cerebellum]. Green bars indicate [internal model AND cerebellum AND motor control], and magenta bars indicate [internal model AND cerebellum AND cognition]. The search was conducted on 6 January 2020.

already too long to be biologically realistic. One of the most difficult tasks in robotics is still bipedal locomotion in rough terrain. Human babies fall only ~ 3000 times during babyhood (Adolph et al., 2012). The three models do not provide insights to explore neural mechanisms for this remarkable capability of the cerebellum and the nervous system in realizing “learning from a small sample” in sensorimotor coordination for systems with huge numbers of degrees of freedom.

Learning of error signals

Second, if it is granted that climbing-fiber inputs provide error signals, and that LTD-LTP of parallel-fiber-Purkinje-cell synapses is the basis of cerebellar supervised learning, the three models do not address how and where learning of error signals originates for higher functions, including manipulation of tools and language acquisition. For phylogenetically old functions, such as the vestibulo-ocular reflex or ocular following responses, error signals carried by climbing-fibers are genetically determined from retinal slips within the accessory optic system. In contrast, tool manipulation or language acquisition is a recent event, occurring within the last several hundred thousand years of human evolution. It is implausible that climbing-fiber error signals are genetically determined for these new functions of the cerebellum. As explained in the Section, Codon theory, the sign and coordinate frame of climbing-fiber error signals should be strictly matched to those of the output from the corresponding part of the deep cerebellar nuclei, so that supervised learning works. The cerebral cortex, which most probably sends original pieces of information to shape climbing-fiber inputs for the new functions, is too far from the

cerebellum to be able to align the sign and coordinates between climbing-fiber input and output from the cerebellum, that is, the cerebellar input–output consistency. The three models do not provide insights about how climbing-fiber inputs are learned in the cerebellum while satisfying cerebellar input–output consistency.

Dimensional mismatch

Third, the three models do not provide theoretical clues to resolve a dimensional mismatch between one degree-of-freedom climbing-fiber inputs and extremely large dimensions to be controlled in whole body movements (Morton and Bastian, 2007; Hoogland et al., 2015; Machado et al., 2015) and higher cognition (Strick et al., 2009; Lu et al., 2012; Schmahmann et al., 2019). Within each microzone, climbing-fiber inputs are similar (Andersson and Oscarsson, 1978; Ito et al., 1982a,b, Apps and Garwicz, 2005; Apps and Hawkes, 2009); thus, the error signal is basically one-dimensional. In oculomotor control, three pairs of extraocular muscles approximately determine horizontal, vertical, and rotational axes of motor coordinates. Accordingly, three microzones of the flocculus possess horizontal, vertical, and rotational retinal slips as climbing-fiber inputs (Andersson and Oscarsson, 1978; Ito et al., 1982a,b). For oculomotor control, the three dimensions of motor coordinates are represented by three microzones of the flocculus. Thus movement and cerebellar dimensions are matched. However, the three models do not provide theoretical insights into how and where extremely large degree-of-freedom systems for whole body motion or cognition can be controlled and learned with single degree-of-freedom climbing-fiber inputs. This theoretical issue is tightly coupled to the first and second difficulties that we explained above. If drastic reduction of dimensions from for example 1000 to only 1 is possible by some neural mechanisms while preserving the sign and coordinate frames of climbing-fiber inputs and cerebellar output so that supervised learning works efficiently, this dimensional reduction simultaneously solves three difficulties: the curse of dimensionality, learning of error signals, and dimensional mismatches between huge problems and cerebellar climbing-fiber inputs. Some recent studies propose possible roles of synchronization of inferior olive neurons by electrical synapses, and meta-cognition, in reduction of effective degree-of-freedom for learning systems to resolve the curse of dimensionality (Tokuda et al., 2010, 2013, 2017; Kawato et al., 2011; Hoang et al., 2020; Cortese et al., 2019a,b).

Recent experimental findings that cannot readily be explained

Recent experimental advances in cerebellar research led to findings that cannot be readily explained by the three models of Marr, Ito, and Albus. These findings include memory formation in granule cells (Yang et al. 2016; Yamada et al., 2019), temporal-difference prediction errors in climbing-fibers (Ohmae and Medina, 2015), reward-related signals in climbing-fibers (Heffley et al.,

2018; Heffley and Hull, 2019; Kostadinov et al., 2019; Larry et al., 2019; Tsutsumi et al., 2019), in parallel fibers (Wagner et al., 2017), and in cerebellar output (Chabrol et al., 2019; Carta et al., 2019; Sendhilnathan et al., 2020), simple-spike dependent memory formation (Lee et al., 2015), control of Purkinje-cell plasticity mediated by molecular-layer interneurons (Rowan et al., 2018), and neuronal heterogeneity between Aldolase C (zebrin) zones (Sillitoe and Joyner 2007; Zhou et al., 2014; Xiao et al. 2014; Tsutsumi et al., 2015, 2019; Tang et al., 2017). Here we postulate hierarchical reinforcement learning with multiple internal models (Kawato et al., 2000; Haruno et al., 2001, 2003; Doya et al., 2002; Sugimoto et al., 2006, 2012a,b; Kawato and Samejima, 2007) as a new computational framework of the cerebellum for resolution of the three theoretical difficulties, as well as to accommodate the above experimental findings (Fig. 5).

Biological constraints for computational exploration

Evolutionarily, the basal ganglia emerged 560 million years ago, the cerebellum 420 million years ago, and the cerebral cortex emerged 250 million years ago. The main goal of the nervous system is to maximize fitness by selecting optimal behaviors. Thus, it is a reinforcement learning task. In specific situations, animals need to achieve appropriate sub-goals that are derived from the main goal, so as to maximize fitness. Sub-goals should include foraging for food, finding mates, and avoiding injury. A fundamental assumption of our hypothesis is that each of the basal ganglia individually, the basal ganglia and the cerebellum combined, or the basal ganglia, the cerebellum, and the cerebral cortex all combined, should be able to execute most important functions at that evolutionary stage, and to add the next most important and essential functions by development and addition of a new brain division (Northcutt, 2002; Shepherd and Rowe, 2017; Naumann et al., 2015; Tosches et al., 2018). We postulate that the three brain divisions, the basal ganglia, the cerebellum, and the cerebral neocortex all have a single computational goal, in contrast to the proposal by Doya (1999). The goal is reinforcement learning to select optimal behaviors, although representations and main computational learning algorithms differ between the three divisions, as discussed in the Section, Synaptic plasticity in achieving supervised learning. The basal ganglia alone in agnathans should provide a plain reinforcement learning system for action selection, with its direct and indirect pathways differentially dependent on dopamine (Ericsson et al., 2011). The plain reinforcement learning system of agnathans is not provided with computational powers of internal models and/or working memory systems that became available only later in evolution. The cerebellum in gnathostomes (fishes with jaws) combined with the basal ganglia should be a reinforcement learning system with hierarchy, modularity, and multiple internal models (Fig. 5). Equipped with 10,000 microzones, enormous numbers of granule cells, and Purkinje-cell LTD-LTP, the cerebellum is well suited for temporal processing of different inputs within several hundred milliseconds

(Medina and Mauk, 2000; Ohmae et al., 2013, 2017; Kunimatsu et al., 2018; Kameda et al., 2019; Sanger and Kawato, 2020). Temporal information processing capability is maximally used in multimodal integration, forward and inverse model learning, and in reward-related information processing (Yamazaki and Lennon, 2019). The cerebral neocortex in mammals combined with the basal ganglia and the cerebellum is an even more sophisticated reinforcement learning system with additional computational powers acquired by pyramidal neuron STDP and abundant recurrent connections. Multimodal integration, associative memory, topographic representations, and cortical temporal dynamics can select abstract, concise, categorical representations of behavioral contexts for fast and efficient reinforcement learning. Those are necessary computations for adaptation and learning of nursing, social and communication behaviors.

Inputs and outputs for reinforcement learning differ in the three brain divisions. While the pyramidal tract contains 2 million-fibers, the globus pallidus internal contain 350,000 neurons (Hardman et al., 2002), and the interpositus nucleus contains only 70,000 neurons in humans (Fukutani et al., 1992; Andersen et al., 2004). Thus, the ratio of output channel bandwidths for direct motor control of the cerebral cortex, the basal ganglia, and the cerebellum is 30:6:1. Consequently, the cerebellum is constrained to act on a set of abstract motor and reward related variables. This is related to the one-dimension error signal spanned by climbing-fiber inputs within a single microzone. If climbing-fiber inputs encode reward/penalty related information, corresponding parts of the cerebellum can learn state-value functions, state-action value functions, reward prediction errors and other reward-related variables from parallel-fiber inputs. The cerebrotocerebellum is connected to the cerebral cortex via both inputs and outputs, and it can utilize efficient, concise spatiotemporal representations acquired in the cerebral cortex (Gao et al., 2018; Wagner et al., 2019). It was recently revealed that for some parts, an output from one microzone is fed to granule cells of another microzone (Houck and Person, 2014; Gao et al., 2016; Giovannucci et al., 2017), implying that these microzones are hierarchically/heterarchically arranged (Ohmae S, private communication) (Fig. 5B). Thus, we have multiple modules, internal models, and hierarchy/heterarchy in the cerebellum.

Hierarchical and modular reinforcement learning in the cerebellum

Hierarchical and/or heterarchical reinforcement learning frameworks with multiple internal models (Kawato et al., 2000; Haruno and Kawato, 2006) could provide theoretical guidelines to resolve the three computational difficulties of the models of Marr, Ito and Albus as well as to interpret new experimental findings. The three computational difficulties of the three models are the curse of dimensionality, learning of error signals, and dimensional mismatch. The proposed frameworks resolve the curse of dimensionality with hierarchical structure because the top layer can search for the optimal solution in a much lower dimension thanks to the dimensional reduction by the bot-




Fig. 5. A hierarchical reinforcement learning model with multiple modules. **(A)** A hierarchical reinforcement learning model with top and bottom layers based on reinforcement MOSAIC (Doya et al., 2002) and hierarchical MOSAIC (Haruno et al., 2003) (adapted from Sugimoto et al., 2006). **(left panel)** The lower curved sheet represents continuous state- x space of the environment with nonlinear dynamics. The continuous space is automatically divided into discrete regions (blue shades) according to goodness of prediction by each forward model, while the MOSAIC architecture predicts the whole nonlinear dynamics with different combinations of forward models, f_i , where the index i is given for each forward model. This discrete division of the continuous state-space allows this MOSAIC to treat the transitions between state-space patches (blue regions) as discrete state transitions in semi-Markov decision processes (i, i', i'' in top layer). **(right panel)** The top layer maximizes a reward r provided by the environment by controlling state- i transition while selecting the abstract action- j , and influencing bottom layer reinforcement learning by providing inter-layer rewards $R_{ij}(x, u)$ for the state i and the action j . In other words, the top layer supplies sub-goal information to the bottom layer. The whole hierarchical reinforcement learning is driven by the reward r , while inter-layer communication was achieved as a state abstraction by forward models in the bottom layer, and setting sub-goals for the bottom layer by the top layer. The top layer symbolically represents which reinforcement-learning modules in the bottom layer are selected, and controls Markov transitions between the discrete states such as i, i', i'' , while selecting the abstract action j that determines inter-layer rewards for the bottom layer. Reinforcement Q-learning takes place with the state-action value function, $Q(i, j)$. $P(j|i)$ denotes j -th action selection probability for the i -th state, i.e., top-layer policy, and is determined from $Q(i, j)$. The bottom layer is a continuous reinforcement learning system to learn an action u such that $R_{ij}(x, u)$ is maximized while the top layer is in the i -th state and takes the action j . When learning is achieved, the top layer maximizes the reward in much smaller dimensions (discrete space of i) than in the original large dimensions of the environment, and the bottom layer maximizes the sub-goals (inter-layer rewards) from the top layer so that the bottom layer faithfully transforms the actions of the top layer in the discrete space into the control of x in the continuous space. $V_{ij}(x)$ is the value function, which is learned from the reward $R_{ij}(x, u)$, bottom-layer state, y , bottom-layer observation, u , bottom-layer action, $f_i(x, u)$, i -th forward model, λ_i^f , responsibility signal of the i -th forward model, u_{ij} is the bottom-layer action under the i -th region with value function $V_{ij}(x)$, and the top-layer state i and action j . Note that there are no inverse models or responsibility predictors in this hierarchical reinforcement MOSAIC. **(B)** Another MOSAIC model with loop structures implements interactions between the three modules for the inverse model, forward model, and responsibility predictor (adapted from Kawato et al., 2000). FM_i , IM_i , and RP_i indicate the i -th forward model, the i -th inverse model, and the i -th responsibility predictor, respectively, realized by Purkinje cells. λ_i is the i -th responsibility signal of the i -th modules, which is common to FM_i , IM_i , and RP_i . x is an actual state. \hat{x}_i is a predicted state by FM_i . $e_i^f = x - \hat{x}_i$ is the prediction error signal of the state, and $\lambda_i e_i^f$ is the final error signal used to change FM_i in supervised learning. \hat{u}_i is the i -th motor command generated by IM_i . $\sum_i \lambda_i \hat{u}_i$ is the responsibility-signal weighted summation of the motor commands from all inverse models, and this is the total motor command output from the cerebellum. u_{fb} is a feedback motor command. $\lambda_i u_{fb}$ is the final error signal to change IM_i used in supervised learning. $\hat{\lambda}_i$ is the i -th predicted responsibility signal computed by RP_i . $\lambda_i - \hat{\lambda}_i$ is the error in responsibility prediction to change RP_i . Red arrows indicate the three different kinds of error signals that drive supervised learning of the three modules. Blue arrows show responsibility signals that control common weighting in supervised learning of the three modules. Brown arrows show prior (predicted responsibility) and likelihood for responsibility estimation. Predicted responsibility signals are computed before movement execution. The three kinds of modules interact with each other in feedback connections between the cerebellar cortex, the dentate nucleus (DE), the red nucleus (RN), and the inferior olive nucleus (IO). VL, ventrolateral thalamus, CBR CX, cerebral cortex, PN, pontine nucleus. *Figure (A) was published in The IEICE transactions on information and systems (Japanese edition), 89(7), Sugimoto N, Samejima K, Doya K, Kawato M, Hierarchical Reinforcement Learning: Temporal Abstraction Based on MOSAIC Model, 1577–1587, Copyright IEICE (2006). *Figure (B) was published in Kagaku, 70(11), Kawato M, Doya K, Haruno M, Multiple paired forward and inverse models (MOSAIC) – the information-processing and possibility, 1009–1017, Copyright Iwanami Shoten (2000).

tom layer and requires only a moderate number of learning trials. Multiple models connected serially rather than in parallel provide a computational architecture where one module can learn the error signal, which is used in learning by the other module. Hierarchical and/or heterarchical structures could connect top and bottom layers with huge differences in dimensions. Here, the top layer may possess very low-dimensional symbolic representations, while the bottom layer may possess extremely high-dimensional representations related to motoneurons and muscles. We note that the three models did not discuss hierarchy, heterarchy, serially connected multiple modules, or how representations with huge differences in dimensionality could be related within the cerebellum. New frameworks may provide several clues to interpret the new experimental findings, and are partially motivated by biological constraints including evolutionary views about the brain.

The hierarchical reinforcement learning scheme with multiple modules shown in Fig. 5A (Sugimoto et al., 2006) illustrates how abstract symbolic representations in the top layer are implemented as responsibility signals in the bottom layer. This model is a combination of a reinforcement MOSAIC (Doya et al., 2002) and hierarchical MOSAIC (Haruno et al., 2003). In each layer, reinforcement learning occurs with multiple modules. The responsibility signal determines which forward model in the bottom layer best represents the environment. Patterns of responsibility signals allow the hierarchical MOSAIC to achieve discretization of continuous space. Changes of dynamics in the continuous space are approximated by Markov transitions between the abstract and discrete states. The top layer controls this transition as well as providing sub-goals to the bottom layer. Drastic dimension reduction is achieved by this abstraction in the top layer. In the bottom layer,




Fig. 5 (continued)

a “divide and conquer” strategy works with many experts (forward models), even for highly nonlinear problems. Abstraction and divide-and-conquer together resolve the curse of dimensionality. In most successful engineering applications of hierarchical reinforcement learning algorithms in robotics, higher-level abstract representations and/or sub-goals of the bottom layer were determined by researchers (Atkeson et al., 2000). So far, intermediate goal postures for a standing robot were manually selected as representations in the top layer by Morimoto and Doya (2001). As another example, Bentivegna et al., (2003) selected right-bank shots as higher-level actions in air-hockey by a humanoid robot DB. Neither the cerebellum nor the whole brain can enjoy the luxurious special treatment that a homunculus in the brain selects appropriate

representations for the top layers. From a neuroscience point of view, abstract representations of movements in the top layers could be movement sub-goals, synergies, motion of center of gravity, locomotion gates, periods and phases of rhythmic movements, and phase relationships of limb rhythms. Computationally, the most difficult question is how appropriate high-level representations can be automatically acquired without homunculus or genetic hard-wiring. Hierarchical and reinforcement MOSAIC can find higher-level representations through the pattern of responsibility signals, that is, which combination of forward models, value functions and policies is most appropriate in a given situation, and could provide a computational possibility for automatically learning sub-goals and synergy.

The heterarchical MOSAIC shown in Fig. 5B (Kawato et al., 2000) shows how the three modules, forward models, inverse models, and responsibility predictors, are heterarchically arranged by convergent and divergent connections within the closed circuit consisting of the dentate, red, and inferior olive nuclei. Responsibility predictors can predict responsibility signals, which estimate goodness of prediction of each forward model from pieces of contextual information. In Allen and Tsukahara (1974) and Kawato and Gomi (1992), functions of the closed loop circuit between the dentate nucleus, the parvo-cellular part of the red nucleus, and the inferior olive

sending climbing-fibers (De Zeeuw et al., 1998; Sokolov et al., 2017) were discussed, but remain enigmatic, although this Ogawa’s triangle (also known as the triangle of Guillaumin and Mollaret) is huge in humans (Ogawa, 1941). A recent study reported that cerebellar output during eyeblink conditioning can contribute to generation of an error signal in climbing-fibers (Ohmae and Medina, 2019). If this triangle is a neural connection from outputs of one microzone to error signals of another, the second theoretical difficulty can be tackled in the hierarchical reinforcement learning framework. In the upper level of hierarchical reinforcement learning, abstract, concise representations could be used and the curse of dimensionality would be drastically relaxed. With the cascade arrangements of microzones, climbing-fiber error signals

for one microzone could be learned by the other upper-layer microzone. Hierarchical structures could bridge concise, abstract representations in the top layer with high-dimensional representations in the bottom layer. One-dimensional error signals for each module in each layer could provide a mechanism to find a low-dimensional order parameter for motor control represented in its outputs.

This is a new era of cerebellar research with veritable explosions of new techniques, new findings, and new computational ideas. It is important and timely to evaluate contributions and limitations of the Marr, Ito, and Albus models and to explore future directions based on advances made following their proposals. In this review, we have made a computationally oriented effort, and in introducing these models we have experienced the same excitement that we felt when we first encountered these theories 50 years ago.

ACKNOWLEDGEMENTS

We thank Drs. Keisuke Toyama, Tomoo Hirano, Soichi Nagao, and Kazuo Kitamura for their comments on an earlier version of the manuscript. This research was partially supported by AMED under Grant Number JP19dm0307008. MK and HH were further supported by JST ERATO (Japan, grant number JPMJER1801).

REFERENCES

- Adolph KE, Cole WG, Komati M, Garciaguire JS, Badaly D, Lingeman JM, Chan GLY, Sotsky RB (2012) How do you learn to walk? Thousands of steps and dozens of falls per day. *Psychol Sci* 23:1387–1394. <https://doi.org/10.1177/0956797612446346>.
- Albus JS (1971) A theory of cerebellar function. *Math Biosci* 10:25–61. [https://doi.org/10.1016/0025-5564\(71\)90051-4](https://doi.org/10.1016/0025-5564(71)90051-4).
- Albus JS (1975) A new approach to manipulator control: the cerebellar model articulation controller (CMAC). *J Dyn Sys Meas Control* 97(3):220–227. <https://doi.org/10.1115/1.3426922>.
- Allen G, Tsukahara N (1974) Cerebrocerebellar communication systems. *Physiol Rev* 54(4):957–1006. <https://doi.org/10.1152/physrev.1974.54.4.957>.
- Amari S (2020) Any target function exists in a neighborhood of any sufficiently wide random network: a geometrical perspective. arXiv:2001.06931.
- Andersen BB, Fabricius K, Gundersen HJG, Jelsing J, Stark AK (2004) No change in neuron numbers in the dentate nucleus of patients with schizophrenia estimated with a new stereological method – the smooth fractionator. *J Anat* 205:313–321. <https://doi.org/10.1111/j.0021-8782.2004.00337.x>.
- Andersson G, Oscarsson O (1978) Projections to lateral vestibular nucleus from cerebellar climbing fiber zones. *Exp Brain Res* 32(4):549–564.
- Anzai M, Kitazawa H, Nagao S (2010) Effects of reversible pharmacological shutdown of cerebellar flocculus on the memory of long-term horizontal vestibulo-ocular reflex adaptation in monkeys. *Neurosci Res* 68(3):191–198. <https://doi.org/10.1016/j.neures.2010.07.2038>.
- Anzai M, Nagao S (2014) Motor learning in common marmosets: vestibulo-ocular reflex adaptation and its sensitivity to inhibitors of Purkinje cell long-term depression. *Neurosci Res* 83:33–42. <https://doi.org/10.1016/j.neures.2014.04.002>.
- Apps R, Garwicz M (2005) Anatomical and physiological foundations of cerebellar information processing. *Nat Rev Neurosci* 6(4):297–311. <https://doi.org/10.1038/nrn1646>.
- Apps R, Hawkes R (2009) Cerebellar cortical organization: a one-map hypothesis. *Nat Rev Neurosci* 10(9):670–681. <https://doi.org/10.1038/nrn2698>.
- Arenz A, Silver RA, Schaefer AT, Margrie TW (2008) The contribution of single synapses to sensory representation in vivo. *Science* 321(5891):977–980. <https://doi.org/10.1126/science.1158391>.
- Armstrong DM, Rawson JA (1979) Activity patterns of cerebellar cortical neurones and climbing fibre afferents in the awake cat. *J Physiol* 289:425–448. <https://doi.org/10.1113/jphysiol.1979.sp012745>.
- Atkeson CG, Benezun PWB, Banerjee N, Berenson D, Bove CP, Cui X, DeDonato M, Du R, et al. (2018) What happened at the DARPA robotics challenge finals. In: Spenko M, Buerger S, Iagnemma K, editors. *The DARPA robotics challenge finals: humanoid robots to the rescue*. Springer Nature. p. 687–1684.
- Atkeson CG, Hale J, Pollick F, Riley M, Kotosaka S, Schaal S, Shibata T, Tevatia G, Vijayakumar S, Ude A, Kawato M (2000) Using humanoid robots to study human behavior. *IEEE Intelligent Syst: Special Issue Humanoid Robotics* 15:46–56.
- Attwell PJ, Cooke SF, Yeo CH (2002) Cerebellar function in consolidation of a motor memory. *Neuron* 34(6):1011–1020. [https://doi.org/10.1016/S0896-6273\(02\)00719-5](https://doi.org/10.1016/S0896-6273(02)00719-5).
- Badura A, De Zeeuw CI (2017) Cerebellar granule cells: dense, rich and evolving representations. *Curr Biol* 27(11):R415–R418. <https://doi.org/10.1016/j.cub.2017.04.009>.
- Barto AG, Sutton RS, Anderson CW (1983) Neuron-like elements that can solve difficult learning control problems. In: *IEEE Trans Syst Man Cybern* 13(5) pp 835–846. IEEE Institute of electrical and electronics engineering, inc.
- Bastian AJ (2006) Learning to predict the future: the cerebellum adapts feedforward movement control. *Curr Opin Neurobiol* 16(6):645–649. <https://doi.org/10.1016/j.conb.2006.08.016>.
- Bell CC, Russell CJ (1978) Effect of electric organ discharge on ampullary receptors in a mormyrid. *Brain Res* 145:85–96. [https://doi.org/10.1016/0006-8993\(78\)90798-9](https://doi.org/10.1016/0006-8993(78)90798-9).
- Bemporad A, Morari M, Dua V, Pistikopoulos EN (2002) The explicit linear quadratic regulator for constrained systems. *Automatica* 38:3–20. [https://doi.org/10.1016/S0005-1098\(01\)00174-1](https://doi.org/10.1016/S0005-1098(01)00174-1).
- Bengtsson F, Jörntell H (2009) Sensory transmission in cerebellar granule cells relies on similarly coded mossy fiber inputs. *Proc Natl Acad Sci U S A* 106(7):2389–2394. <https://doi.org/10.1073/pnas.0808428106>.
- Bentivegna DC, Atkeson CG, Cheng G (2003) Learning to select primitives and generate sub-goals from practice. *Intelligent Robots Systems* 1:946–953. <https://doi.org/10.1109/IROS.2003.1250750>.
- Bhanpuri NH, Okamura AM, Bastian AJ (2014) Predicting and correcting ataxia using a model of cerebellar function. *Brain* 137(7):1931–1944. <https://doi.org/10.1093/brain/awu115>.
- Bhushan N, Shadmehr R (1999) Computational nature of human adaptive control during learning of reaching movements in force fields. *Biol Cybern* 81(1):39–60. <https://doi.org/10.1007/s004220050543>.
- Bi G, Poo M (2001) Synaptic modification by correlated activity: Hebb's postulate revisited. *Annu Rev Neurosci* 24:139–166. <https://doi.org/10.1146/annurev.neuro.24.1.139>.
- Blakemore SJ, Frith CD, Wolpert DM (2001) The cerebellum is involved in predicting the sensory consequences of action. *NeuroReport* 12:1879–1884. <https://doi.org/10.1097/00001756-200107030-00023>.
- Blakemore SJ, Wolpert DM, Frith CD (1998) Central cancellation of self-produced tickle sensation. *Nat Neurosci* 1:635–640. <https://doi.org/10.1038/2870>.
- Brindley GS (1964) The use made by the cerebellum of the information that it receives from sense organs. *Int Brain Res Org Bulletin* 3:80.
- Brindley GS (1969) Nerve net models of plausible size that perform many simple learning tasks. *Proc R Soc Lond* 174(1035):173–191. <https://doi.org/10.1098/rspb.1969.0087>.
- Brooks JX, Carriot J, Cullen KE (2015) Learning to expect the unexpected: rapid updating in primate cerebellum during voluntary self-motion. *Nat Neurosci* 18:1310–1317. <https://doi.org/10.1038/nn.4077>.

- Brooks JX, Cullen KE (2013) The primate cerebellum selectively encodes unexpected self-motion. *Curr Biol* 23(11):947–955. <https://doi.org/10.1016/j.cub.2013.04.029>.
- Caporale N, Dan Y (2008) Spike timing-dependent plasticity: a Hebbian learning rule. *Annu Rev Neurosci* 31:25–46. <https://doi.org/10.1146/annurev.neuro.31.060407.125639>.
- Carta I, Chen CH, Schott A, Dorizan S, Khodakhah K (2019) Cerebellar modulation of the reward circuitry and social behavior. *Science* 363(6424):eaav0581. <https://doi.org/10.1126/science.aav0581>.
- Chabrol FP, Arenz A, Wiechert MT, Margrie TW, DiGregorio DA (2015) Synaptic diversity enables temporal coding of coincident multi-sensory inputs in single neurons. *Nat Neurosci* 18(5):718–727. <https://doi.org/10.1038/nn.3974>.
- Chabrol FP, Blot A, Mrsic-Flogel TD (2019) Cerebellar contribution to preparatory activity in motor neocortex. *Neuron* 103:506–519. <https://doi.org/10.1016/j.neuron.2019.05.022>.
- Chaumont J, Guyon N, Valera AM, Dugué GP, Popa D, Marcaggi P, Gautheron V, Reibel-Foisset S, et al. (2013) Clusters of cerebellar Purkinje cells control their afferent climbing fiber discharge. *Proc Natl Acad Sci U S A* 110(40):16223–16228. <https://doi.org/10.1073/pnas.1302310110>.
- Cortese A, De Martino B, Kawato M (2019) The neural and cognitive architecture for learning from a small sample. *Curr Opin Neurobiol* 55:133–214. <https://doi.org/10.1016/j.conb.2019.02.011>.
- Cortese A, Lau H, Kawato M (2019b) Metacognition facilitates the exploitation of unconscious brain states. *bioRxiv* doi:10.1101/548941
- Cover TM (1965) Geometrical and statistical properties of systems of linear inequalities with applications in pattern recognition. *IEEE Trans Elec Comput* 3:326–334. <https://doi.org/10.1109/PGEC.1965.264137>.
- Cybenko G (1989) Approximation by superpositions of a sigmoidal function. *Math Cont Sig Syst* 2:303–314. <https://doi.org/10.1007/BF02551274>.
- D'Angelo E, Mapelli L, Casellato C, Garrido JA, Luque N, Monaco J, Prestori F, Pedrocchi A, et al. (2016) Distributed circuit plasticity: new clues for the cerebellar mechanisms of learning. *Cerebellum* 15(2):139–151. <https://doi.org/10.1007/s12311-015-0711-7>.
- Deverett B, Kislin M, Tank DW, Wang SS (2019) Cerebellar disruption impairs working memory during evidence accumulation. *Nat Commun* 10:3128. <https://doi.org/10.1038/s41467-019-11050-x>.
- De Zeeuw CI, Simpson JI, Hoogenraad CC, Galjart N, Koekkoek SK, Ruijgrok TJ (1998) Microcircuitry and function of the inferior olive. *Trends Neurosci* 21(9):391–400.
- Doi T, Kuroda S, Michikawa T, Kawato M (2005) Inositol 1,4,5-trisphosphate-dependent Ca²⁺ + threshold dynamics detect spike timing in cerebellar Purkinje cells. *J Neurosci* 25:950–961. <https://doi.org/10.1523/JNEUROSCI.2727-04.2005>.
- Doya K (1999) What are the computations of the cerebellum, the basal ganglia and the cerebral cortex? *Neural Netw* 12(7–8):961–974. [https://doi.org/10.1016/S0893-6080\(99\)00046-5](https://doi.org/10.1016/S0893-6080(99)00046-5).
- Doya K, Samejima K, Katagiri K, Kawato M (2002) Multiple model-based reinforcement learning. *Neural Computation* 14:1347–1369. <https://doi.org/10.1162/089976602753712972>.
- Eccles J, Ito M, Szentagothai J (1967) The cerebellum as a neuronal machine. Springer science + business media New York.
- El-Shamayleh Y, Kojima Y, Soetedjo R, Horwitz GD (2017) Selective optogenetic control of Purkinje cells in monkey cerebellum. *Neuron* 95(1):51–62.e4. <https://doi.org/10.1016/j.neuron.2017.06.002>.
- Ericsson J, Silberberg G, Robertson B, Wikström MA, Grillner S (2011) Striatal cellular properties conserved from lampreys to mammals. *J Physiol* 589(Pt 12):2979–2992. <https://doi.org/10.1113/jphysiol.2011.209643>.
- Fernandez E, Schiappa R, Girault JA, Le Novere N (2006) DARPP-32 is a robust integrator of dopamine and glutamate signals. *PLoS Comput Biol* 2. <https://doi.org/10.1371/journal.pcbi.0020176> e176.
- Fujita M (1982) Adaptive filter model of the cerebellum. *Biol Cybern* 45(3):195–206. <https://doi.org/10.1007/bf00336192>.
- Fukutani Y, Nakamura I, Kobayashi K, Yamaguchi N, Matsubara R (1992) Pathology of the cerebellar dentate and interpositus nuclei in Joseph disease: a morphometric investigation. *J Neurol Sci* 108:39–47. [https://doi.org/10.1016/0022-510X\(92\)90186-O](https://doi.org/10.1016/0022-510X(92)90186-O).
- Funahashi K (1989) On the approximate realization of continuous mappings by neural networks. *Neural Networks* 2(3):183–192. [https://doi.org/10.1016/0893-6080\(89\)90003-8](https://doi.org/10.1016/0893-6080(89)90003-8).
- Gao Z, Davis C, Thomas AM, Economo MN, Abrego AM, Svoboda K, De Zeeuw CI, Li N (2018) A cortico-cerebellar loop for motor planning. *Nature* 563(7729):113–116. <https://doi.org/10.1038/s41586-018-0633-x>.
- Gao Z, Proietti-Onori M, Lin Z, Ten Brinke MM, Boele HJ, Potters JW, Ruijgrok TJ, Hoebeek FE, et al. (2016) Excitatory cerebellar nucleocortical circuit provides internal amplification during associative conditioning. *Neuron* 89(3):645–657. <https://doi.org/10.1016/j.neuron.2016.01.008>.
- Gao Z, Van Beugen BJ, De Zeeuw CI (2012) Distributed synergistic plasticity and cerebellar learning. *Natu Rev Neurosci* 13(9):619–635. <https://doi.org/10.1038/nrn3312>.
- Gardner E, Derrida B (1988) Optimal storage properties of neural network models. *J Phys A* 21(1):271. <https://doi.org/10.1088/0305-4470/21/1/031>.
- Gilmer JI, Person AL (2018) Theoretically sparse, empirically dense: new views on cerebellar granule cells. *Trends Neurosci* 41(12):874–877. <https://doi.org/10.1016/j.tins.2018.09.013>.
- Giovannucci A, Badura A, Deverett B, Najafi F, Pereira TD, Gao Z, Ozden I, Kloth AD, et al. (2017) Cerebellar granule cells acquire a widespread predictive feedback signal during motor learning. *Nat Neurosci* 20(5):727–734. <https://doi.org/10.1038/nn.4531>.
- Gomi H, Kawato M (1992) Adaptive feedback control models of the vestibulocerebellum and spinocerebellum. *Biological Cybernetics* 68:105–114. <https://doi.org/10.1007/BF00201432>.
- Gomi H, Kawato M (1993) Recognition of manipulated objects by motor learning with modular architecture networks. *Neural Networks* 6:485–497. [https://doi.org/10.1016/S0893-6080\(05\)80053-X](https://doi.org/10.1016/S0893-6080(05)80053-X).
- Gomi H, Kawato M (1996) Equilibrium-point control hypothesis examined by measured arm stiffness during multijoint movement. *Science* 272(5258):117–120. <https://doi.org/10.1126/science.272.5258.117>.
- Gomi H, Shidara M, Takemura A, Inoue Y, Kawano K, Kawato M (1998) Temporal firing patterns of Purkinje cells in the cerebellar ventral paraflocculus during ocular following responses in monkeys I. Simple spikes. *J Neurophysiol* 80(2):818–831. <https://doi.org/10.1152/jn.1998.80.2.818>.
- Han VZ, Grant G, Bell CC (2000) Reversible associative depression and nonassociative potentiation at a parallel fiber synapse. *Neuron* 27:611–622. [https://doi.org/10.1016/S0896-6273\(00\)00070-2](https://doi.org/10.1016/S0896-6273(00)00070-2).
- Hansel C, Linden DJ, D'Angelo E (2001) Beyond parallel fiber LTD: the diversity of synaptic and non-synaptic plasticity in the cerebellum. *Nat Neurosci* 4(5):467–475. <https://doi.org/10.1038/87419>.
- Hardman CD, Henderson JM, Finkelstein DI, Horne MK, Paxinos G, Halliday GM (2002) Comparisons of the basal ganglia in rats, marmosets, macaques, baboons, and humans: volume and neuronal number for the output, internal relay, and striatal modulating nuclei. *J Comp Neurol* 445:238–255. <https://doi.org/10.1002/cne.10165>.
- Haruno M, Kawato M (2006) Heterarchical reinforcement-learning model for integration of multiple cortico-striatal loops; fMRI examination in stimulus-action-reward association learning. *Neural Networks* 19:1242–1254.
- Haruno M, Wolpert DM, Kawato M (2001) MOSAIC model for sensorimotor learning and control. *Neural Comput* 13:2201–2220. <https://doi.org/10.1162/089976601750541778>.
- Haruno M, Wolpert DM, Kawato M (2003) Hierarchical MOSAIC for movement generation. *Int Congress Series* 1250:575–590.

- Heffley W, Hull C (2019) Classical conditioning drives learned reward prediction signals in climbing fibers across the lateral cerebellum. *eLife* 8: e46764. doi: 10.7554/eLife.46764.001.
- Heffley W, Song EY, Xu Z, Taylor BN, Hughes MA, McKinney A, Joshua M, Hull C (2018) Coordinated cerebellar climbing fiber activity signals learned sensorimotor predictions. *Nat Neurosci* 21(10):1431–1441. <https://doi.org/10.1038/s41593-018-0228-8>.
- Heiney SA, Kim J, Augustine GJ, Medina JF (2014) Precise control of movement kinematics by optogenetic inhibition of Purkinje cell activity. *J Neurosci* 34(6):2321–2330. <https://doi.org/10.1523/JNEUROSCI.4547-13.2014>.
- Herzfeld DJ, Kojima Y, Soetedjo R, Shadmehr R (2015) Encoding of action by the Purkinje cells of the cerebellum. *Nature* 526(7573):439–442. <https://doi.org/10.1038/nature15693>.
- Herzfeld DJ, Kojima Y, Soetedjo R, Shadmehr R (2018) Encoding of error and learning to correct that error by the Purkinje cells of the cerebellum. *Nat Neurosci* 21(5):736–743. <https://doi.org/10.1038/s41593-018-0136-y>.
- Higuchi S, Chaminade T, Imamizu H, Kawato M (2009) Shared neural correlates for language and tool-use in Broca's area. *NeuroReport* 20:1376–1381. <https://doi.org/10.1097/WNR.0b013e3283315570>.
- Higuchi S, Imamizu H, Kawato M (2007) Cerebellar activity evoked by common tool-use execution and imagery tasks: an fMRI study. *Cortex* 3:350–358. [https://doi.org/10.1016/S0010-9452\(08\)70460-X](https://doi.org/10.1016/S0010-9452(08)70460-X).
- Hirano T (1990) Depression and potentiation of the synaptic transmission between a granule cell and a Purkinje cell in rat cerebellar culture. *Neurosci Lett* 119:141–144. [https://doi.org/10.1016/0304-3940\(90\)90818-T](https://doi.org/10.1016/0304-3940(90)90818-T).
- Hirano T (2013) Long-term depression and other synaptic plasticity in the cerebellum. *Proc Jpn Acad Ser B Phys Biol Sci* 89(5):183–195. <https://doi.org/10.2183/pjab.89.183>.
- Hoang H, Lang EJ, Hirata Y, Tokuda IT, Aihara K, Toyama K, Kawato M & Schweighofer N (2020): Electrical coupling controls dimensionality and chaotic firing of inferior olive neurons, *PLoS Computational Biology*, in press.
- Honda T, Nagao S, Hashimoto Y, Ishikawa K, Yokota T, Mizusawa H, Ito M (2018) Tandem internal models execute motor learning in the cerebellum. *Proc Natl Acad Sci USA* 115:7428–7433.
- Hoogland TM, De Groot JR, Witter L, Canto CB, De Zeeuw CI (2015) Role of synchronous activation of cerebellar Purkinje cell ensembles in multi-joint movement control. *Curr Biol* 25(9):1157–1165. <https://doi.org/10.1016/j.cub.2015.03.009>.
- Hopfield JJ (1982) Neural networks and physical systems with emergent collective computational properties. *Proc Nat Acad Sci (USA)* 79:2554–2558. <https://doi.org/10.1073/pnas.79.8.2554>.
- Houck BD, Person AL (2014) Cerebellar loops: a review of the nucleocortical pathway. *Cerebellum* 13(3):378–385. <https://doi.org/10.1007/s12311-013-0543-2>.
- Huang CC, Sugino K, Shima Y, Guo C, Bai S, Mensh BD, Nelson SB, Hantman AW (2013) Convergence of pontine and proprioceptive streams onto multimodal cerebellar granule cells. *eLife* 2. <https://doi.org/10.7554/eLife.00400> e00400.
- Imamizu H, Kuroda T, Miyauchi S, Yoshioka T, Kawato M (2003) Modular organization of internal models of tools in the human cerebellum. *Proc Natl Acad Sci USA* 100:5461–5466. <https://doi.org/10.1073/pnas.0835746100>.
- Imamizu H, Miyauchi S, Tamada T, Sasaki Y, Takino R, Puetz B, Yoshioka T, Kawato M (2000) Human cerebellar activity reflecting an acquired internal model of a new tool. *Nature* 403:192–195. <https://doi.org/10.1038/35003194>.
- Inoshita T, Hirano T (2018) Occurrence of long-term depression in the cerebellar flocculus during adaptation of optokinetic response. *eLife*, 7:36209. doi: 10.7554/eLife.36209.
- Inoue M, Kitazawa S (2018) Motor error in parietal area 5 and target error in area 7 drive distinctive adaptation in reaching. *Curr Biol* 28(14):2250–2262.e3. <https://doi.org/10.1016/j.cub.2018.05.056>.
- Inoue M, Uchimura M, Kitazawa S (2016) Error signals in motor cortices drive adaptation in reaching. *Neuron* 90(5):1114–1126. <https://doi.org/10.1016/j.neuron.2016.04.029>.
- Ishikawa T, Shimuta M, Häusser M (2015). Multimodal sensory integration in single cerebellar granule cells in vivo. *eLife* 4: e12916. doi: 10.7554/eLife.12916.
- Ito M (1970) Neurophysiological aspects of the cerebellar motor control system. *Int J Neurol* 7(2):162–176.
- Ito M (1974) The control mechanisms of cerebellar motor systems. In: Schmitt FO, Worden FG, editors. *The neurosciences, third study program*. MIT Press. p. 293–303.
- Ito M (1984) *The Cerebellum and Neural Control* (Ito M) Raven Pr
- Ito M (2001) Cerebellar long-term depression: characterization, signal transduction, and functional roles. *Physiol Rev* 81(3):1143–1195. <https://doi.org/10.1152/physrev.2001.81.3.1143>.
- Ito M (2002) The molecular organization of cerebellar long-term depression. *Nat Rev Neurosci* 3:896–902. <https://doi.org/10.1038/nrn962>.
- Ito M (2006) Cerebellar circuitry as a neuronal machine. *Prog Neurobiol* 78:272–303. <https://doi.org/10.1016/j.pneurobio.2006.02.006>.
- Ito M (2008) Control of mental activities by internal models in the cerebellum. *Nat Rev Neurosci* 9(4):304–313. <https://doi.org/10.1038/nrn2332>.
- Ito M (2011) *The Cerebellum: Brain for an Implicit Self* (FT Press Science) 1st Edition (Ito M) FT Press.
- Ito M, Kano M (1982) Long-lasting depression of parallel fiber-Purkinje cell transmission induced by conjunctive stimulation of parallel fibers and climbing fibers in the cerebellar cortex. *Neurosci Lett* 33(3):253–258. [https://doi.org/10.1016/0304-3940\(82\)90380-9](https://doi.org/10.1016/0304-3940(82)90380-9).
- Ito M, Orlov I, Yamamoto M (1982a) Topographical representation of vestibulo-ocular reflexes in rabbit cerebellar flocculus. *Neuroscience* 7(7):1657–1664. [https://doi.org/10.1016/0306-4522\(82\)90024-0](https://doi.org/10.1016/0306-4522(82)90024-0).
- Ito M, Sakurai M, Tongroach P (1982b) Climbing fibre induced depression of both mossy fibre responsiveness and glutamate sensitivity of cerebellar Purkinje cells. *J Physiol* 324:113–134. <https://doi.org/10.1113/jphysiol.1982.sp014103>.
- Ito M, Yamaguchi K, Nagao S, Yamazaki T (2014) Long-term depression as a model of cerebellar plasticity. *Prog Brain Res* 210:1–30. <https://doi.org/10.1016/B978-0-444-63356-9.00001-7>.
- Ito M, Yoshida M, Obata K (1964) Monosynaptic inhibition of the intracerebellar nuclei induced from the cerebellar cortex. *Experientia* 20:575–576. <https://doi.org/10.1007/BF02150304>.
- Ito M, Yoshida M (1964) The cerebellar-evoked monosynaptic inhibition of Deiters' neurones. *Experientia* 20(9):515–516. <https://doi.org/10.1007/bf02154085>.
- Izawa J, Criscimagna-Hemminger SE, Shadmehr (2012) Cerebellar contributions to reach adaptation and learning sensory consequences of action. *J Neurosci* 32(12):4230–4239. <https://doi.org/10.1523/JNEUROSCI.6353-11.2012>.
- Jacobs RA, Jordan MI, Nowlan SJ, Hinton GE (1991) Adaptive mixtures of local experts. *Neural Comput* 3(1):79–87. <https://doi.org/10.1162/neco.1991.3.1.79>.
- Jordan MI, Rumelhart DE (1992) Forward models: supervised learning with a distal teacher. *Cognitive Sci* 16(3):307–354. [https://doi.org/10.1016/0364-0213\(92\)90036-T](https://doi.org/10.1016/0364-0213(92)90036-T).
- Jörntell H, Ekerot CF (2006) Properties of somatosensory synaptic integration in cerebellar granule cells in vivo. *J Neurosci* 26(45):11786–11797. <https://doi.org/10.1523/JNEUROSCI.2939-06.2006>.
- Kakegawa W, Katoh A, Narumi S, Miura E, Motohashi J, Takahashi A, Kohda K, Fukazawa Y, et al. (2018) Optogenetic control of synaptic AMPA receptor endocytosis reveals roles of LTD in motor learning. *Neuron* 99(5):985–998.e6. <https://doi.org/10.1016/j.neuron.2018.07.034>.
- Kameda M, Ohmae S, Tanaka M. (2019) Entrained neuronal activity to periodic visual stimuli in the primate striatum compared with the cerebellum. *eLife* 8:e48702. doi: 10.7554/eLife.48702.
- Kawano K (1999) Ocular tracking: behavior and neurophysiology. *Curr Opin Neurobiol* 9(4):467–473. [https://doi.org/10.1016/S0959-4388\(99\)80070-1](https://doi.org/10.1016/S0959-4388(99)80070-1).

- Kawano K, Shidara M (1993) The role of the ventral paraflocculus in ocular following in the monkey. In: Mano N, Hamada I, DeLong MR, editors. *Role of the cerebellum and basal ganglia in voluntary movement*. Excerpta Medica. p. 195–202.
- Kawano K, Shidara M, Takemura A, Inoue Y, Gomi H, Kawato M (1996a) Inverse-dynamics representation of eye movements by cerebellar Purkinje cell activity during short-latency ocular-following responses. *Ann N Y Acad Sci* 781:314–321. <https://doi.org/10.1111/j.1749-6632.1996.tb15709.x>.
- Kawano K, Takemura A, Inoue Y, Kitama T, Kobayashi Y, Mustari MJ (1996b) Visual inputs to cerebellar ventral paraflocculus during ocular following responses. *Prog Brain Res* 112:415–422. [https://doi.org/10.1016/s0079-6123\(08\)63346-4](https://doi.org/10.1016/s0079-6123(08)63346-4).
- Kawato M (1999) Internal models for motor control and trajectory planning. *Curr Opin Neurobiol* 9(6):718–727. [https://doi.org/10.1016/s0959-4388\(99\)00028-8](https://doi.org/10.1016/s0959-4388(99)00028-8).
- Kawato M, Doya K, Haruno M (2000) Multiple paired forward and inverse models (MOSAIC) - information processing and possibility. *Kagaku* 70(11):1009–1017.
- Kawato M, Furukawa K, Suzuki R (1987) A hierarchical neural-network model for control and learning of voluntary movement. *Biol Cybern* 57(3):169–185. <https://doi.org/10.1007/BF00364149>.
- Kawato M, Gomi H (1992) A computational model of four regions of the cerebellum based on feedback-error learning. *Biol Cybernetics* 68:95–103. <https://doi.org/10.1007/BF00201431>.
- Kawato M, Kuroda S, Schweighofer N (2011) Cerebellar supervised learning revisited: biophysical modeling and degrees-of-freedom control. *Curr Opin Neurobiol* 21(5):791–800. <https://doi.org/10.1016/j.conb.2011.05.014>.
- Kawato M, Kuroda T, Imamizu H, Nakano E, Miyauchi S, Yoshioka T (2003) Internal forward models in the cerebellum: fMRI study on grip force and load force coupling. In: Prablanc C, Pelisson D, Rossetti Y, editors. *Progress in Brain Research* 142 "Neural Control of Space Coding and Action Production. The Netherlands: Elsevier, Amsterdam. p. 171–188. doi: 10.1016/S0079-6123(03)42013-X.
- Kawato M, Samejima K (2007) Efficient reinforcement learning: computational theories, neuroscience and robotics. *Curr Opin Neurobiol* 17:205–212. <https://doi.org/10.1016/j.conb.2007.03.004>.
- Kennedy A, Wayne G, Kaifosh P, Alvina K, Abbott LF, Sawtell NB (2014) A temporal basis for predicting the sensory consequences of motor commands in an electric fish. *Nat Neurosci* 17(3):416–422. <https://doi.org/10.1038/nn.3650>.
- Keser Z, Hasan KM, Mwangi BI, Kamali A, Ucisik-Keser FE, Riascos RF, Yozbatiran N, Francisco GE, et al. (2015) Diffusion tensor imaging of the human cerebellar pathways and their interplay with cerebral macrostructure. *Front Neuroanat* 9:41. <https://doi.org/10.3389/fnana.2015.00041>.
- Khilkevich A, Canton-Josh J, DeLord E, Mauk MD (2018) A cerebellar adaptation to uncertain inputs. *Sci Adv* 4(5):eaap9660. <https://doi.org/10.1126/sciadv.aap9660>.
- Kitazawa S, Kimura T, Yin PB (1998) Cerebellar complex spikes encode both destinations and errors in arm movements. *Nature* 392(6675):494–497. <https://doi.org/10.1038/33141>.
- Knogler LD, Markov DA, Dragomir EI, Štíh V, Portugues R (2017) Sensorimotor representations in cerebellar granule cells in larval zebrafish are dense, spatially organized, and non-temporally patterned. *Curr Biol* 27(9):1288–1302. <https://doi.org/10.1016/j.cub.2017.03.029>.
- Kobayashi Y, Kawano K, Takemura A, Inoue Y, Kitama T, Gomi H, Kawato M (1998) Temporal firing patterns of Purkinje cells in the cerebellar ventral paraflocculus during ocular following responses in monkeys. II. complex spikes. *J Neurophysiol* 80:832–848. <https://doi.org/10.1152/jn.1998.80.2.832>.
- Kono M, Kakegawa W, Yoshida K, Yuzaki M (2019) Interneuronal NMDA receptors regulate long-term depression and motor learning in the cerebellum. *J Physiol* 597(3):903–920. <https://doi.org/10.1113/JP276794>.
- Kostadinov D, Beau M, Blanco-Pozo M, Häusser M (2019) Predictive and reactive reward signals conveyed by climbing fiber inputs to cerebellar Purkinje cells. *Nat Neurosci* 22(6):950–962. <https://doi.org/10.1038/s41593-019-0381-8>.
- Kotaleski JH, Lester D, Blackwell KT (2002) Subcellular interactions between parallel fibre and climbing fibre signals in Purkinje cells predict sensitivity of classical conditioning to interstimulus interval. *Integr Physiol Behav Sci* 37:265–292. <https://doi.org/10.1007/bf02734249>.
- Kunimatsu J, Suzuki TW, Ohmae S, Tanaka M. (2018) Different contributions of preparatory activity in the basal ganglia and cerebellum for self-timing. *eLife* 7:e35676. doi: 10.7554/eLife.35676.
- Kuroda S, Schweighofer N, Kawato M (2001) Exploration of signal transduction pathways in cerebellar long-term depression by kinetic simulation. *J Neurosci* 21:5693–5702. <https://doi.org/10.1523/JNEUROSCI.21-15-05693.2001>.
- Larry N, Yarkoni M, Lixenberg A, Joshua M (2019) Cerebellar climbing fibers encode expected reward size. *eLife* 8:e46870. doi: 10.7554/eLife.46870.
- Laurens J, Meng H, Angelaki DE (2013) Neural representation of orientation relative to gravity in the macaque cerebellum. *Neuron* 80(6):1508–1518. <https://doi.org/10.1016/j.neuron.2013.09.029>.
- Lee KH, Mathews PJ, Reeves AMB, Choe KY, Jami SA, Serrano RE, Otis TS (2015) Circuit mechanisms underlying motor memory formation in the cerebellum. *Neuron* 86(2):529–540. <https://doi.org/10.1016/j.neuron.2015.03.010>.
- Li W, Todorov E (2004) Iterative linear quadratic regulator design for nonlinear biological movement systems. In: *Proceedings of the 1st International Conference on Informatics in Control, Automation and Robotics*. pp 222–229.
- Linden DJ (1999) The return of the spike: postsynaptic action potentials and the induction of LTP and LTD. *Neuron* 22:661–666. [https://doi.org/10.1016/s0896-6273\(00\)80726-6](https://doi.org/10.1016/s0896-6273(00)80726-6).
- Linden DJ, Connor JA (1995) Long-term synaptic depression. *Annu Rev Neurosci* 18:319–357. <https://doi.org/10.1146/annurev.ne.18.030195.001535>.
- Lindskog M, Kim M, Wikstrom MA, Blackwell KT, Kotaleski JH (2006) Transient calcium and dopamine increase PKA activity and DARPP-32 phosphorylation. *PLoS Comput Biol* 2. <https://doi.org/10.1371/journal.pcbi.0020119> e119.
- Lisberger SG, Sejnowski TJ (1992) Motor learning in a recurrent network model based on the vestibulo-ocular reflex. *Nature* 360(6400):159–161. <https://doi.org/10.1038/360159a0>.
- Llinas R, Sugimori M (1980) Electrophysiological properties of in vitro Purkinje cell dendrites in mammalian cerebellar slices. *J Physiol* 305:197–213. <https://doi.org/10.1113/jphysiol.1980.sp013358>.
- Lu X, Miyachi S, Takada M (2012) Anatomical evidence for the involvement of medial cerebellar output from the interpositus nuclei in cognitive functions. *Proc Natl Acad Sci U S A* 109(46):18980–18984. <https://doi.org/10.1073/pnas.1211168109>.
- Machado AS, Darmohray DM, Fayad J, Marques HG, Carey MR (2015) A quantitative framework for whole-body coordination reveals specific deficits in freely walking ataxic mice. *eLife* 4. <https://doi.org/10.7554/eLife.07892> e07892.
- Markram H, Lubke J, Frotscher M, Sakmann B (1997) Regulation of synaptic efficacy by coincidence of postsynaptic APs and EPSPs. *Science* 275:213–215. <https://doi.org/10.1126/science.275.5297.213>.
- Marr D (1969) A theory of cerebellar cortex. *J Physiol (Lond)* 202:437–470. <https://doi.org/10.1113/jphysiol.1969.sp008820>.
- Marr D (1982) *Vision: A Computational Investigation into the Human Representation and Processing of Visual Information* (Marr D) W H Freeman & Co.
- Maruta J, Hensbroek RA, Simpson JI (2007) Intra-burst and inter-burst signaling by climbing fibers. *J Neurosci* 27(42):11263–11270. <https://doi.org/10.1523/JNEUROSCI.2559-07.2007>.
- Mathy A, Ho SS, Davie JT, Duguid IC, Clark BA, Häusser M (2009) Encoding of oscillations by axonal bursts in inferior olive neurons. *Neuron* 62(3):388–399. <https://doi.org/10.1016/j.neuron.2009.03.023>.

- Medina JF, Garcia KS, Mauk MD (2001) A mechanism for savings in the cerebellum. *J Neurosci* 21(11):4081–4089. <https://doi.org/10.1523/JNEUROSCI.21-11-04081.2001>.
- Medina JF, Mauk MD (2000) Computer simulation of cerebellar information processing. *Nat Neurosci* 3(Suppl):1205–1211. <https://doi.org/10.1038/81486>.
- Medina JF, Repa JC, Mauk MD, LeDoux JE (2002) Parallels between cerebellum- and amygdala-dependent conditioning. *Nat Rev Neurosci* 3:122–131. <https://doi.org/10.1038/nrn728>.
- Mehta B, Schaal S (2002) Forward models in visuomotor control. *J Neurophysiol* 88(2):942–953.
- Miall RC, Weir DJ, Wolpert DM, Stein JF (1993) Is the cerebellum a smith predictor? *J Mot Behav* 25(3):203–216. <https://doi.org/10.1080/00222895.1993.9942050>.
- Miles FA, Lisberger SG (1981) Plasticity in the vestibulo-ocular reflex: a new hypothesis. *Annu Rev Neurosci* 4:273–299. <https://doi.org/10.1146/annurev.ne.04.030181.001421>.
- Miyamoto H, Kawato M, Setoyama T, Suzuki R (1988) Feedback-error-learning neural network for trajectory control of a robotic manipulator. *Neural Networks* 1:251–265. [https://doi.org/10.1016/0893-6080\(88\)90030-5](https://doi.org/10.1016/0893-6080(88)90030-5).
- Morimoto J, Doya K (2001) Acquisition of stand-up behavior by a real robot using hierarchical reinforcement learning. *Robotics Autonomous Syst* 36:37–51. [https://doi.org/10.1016/S0921-8890\(01\)00113-0](https://doi.org/10.1016/S0921-8890(01)00113-0).
- Morton SM, Bastian AJ (2007) Mechanisms of cerebellar gait ataxia. *Cerebellum* 6(1):79–86. <https://doi.org/10.1080/14734220601187741>.
- Nakano T, Doi T, Yoshimoto J, Doya K (2010) A kinetic model of dopamine- and calcium-dependent striatal synaptic plasticity. *PLoS Comput Biol* 6. <https://doi.org/10.1371/journal.pcbi.1000670> e1000670.
- Naumann RK, Ondracek JM, Reiter S, Shein-Idelson M, Tosches MA, Yamawaki TM, Laurent G (2015) The reptilian brain. *Curr Biol* 25(8):R317–R322. <https://doi.org/10.1016/j.cub.2015.02.049>.
- Nicholson DA, Freeman Jr JH (2003) Addition of inhibition in the olivocerebellar system and the ontogeny of a motor memory. *Nat Neurosci* 6(5):532–537. <https://doi.org/10.1038/nn1042>.
- Northcutt RG (2002) Understanding vertebrate brain evolution. *Integr Comp Biol* 42(4):743–756. <https://doi.org/10.1093/icb/42.4.743>.
- Obata K, Ito M, Ochi R, Sato N (1967) Pharmacological properties of the postsynaptic inhibition by Purkinje cell axons and the action of γ -aminobutyric acid on deiters neurons. *Exp Brain Res* 4:43–57.
- Ogasawara H, Doi T, Doya K, Kawato M (2007) Nitric oxide regulates input specificity of long-term depression and context dependence of cerebellar learning. *PLoS Comput Biol* 3(1). <https://doi.org/10.1371/journal.pcbi.0020179> e179.
- Ogasawara H, Doi T, Kawato M (2008) Systems biology perspectives on cerebellar long-term depression. *NeuroSignals* 16:300–317. <https://doi.org/10.1159/000123040>.
- Ogawa T (1941) Ein Dreieck: nucleus ruber - nucleus olivae - nucleus dentatus. *Jpn J Med Sci* 9:57–65.
- Ohmae S, Uematsu A, Tanaka M (2013) Temporally specific sensory signals for the detection of stimulus omission in the primate deep cerebellar nuclei. *J Neurosci* 33(39):15432–15441. <https://doi.org/10.1523/JNEUROSCI.1698-13.2013>.
- Ohmae S, Kunimatsu J, Tanaka M (2017) Cerebellar roles in self-timing for sub- and supra-second intervals. *J Neurosci* 37(13):3511–3522. <https://doi.org/10.1523/JNEUROSCI.2221-16.2017>.
- Ohmae S, Medina JF (2015) Climbing fibers encode a temporal-difference prediction error during cerebellar learning in mice. *Nat Neurosci* 18(12):1798–1803. doi: 10.1038/nn.4167.
- Ohmae S, Medina JF (2019) Plasticity of ponto-cerebellar circuits generates a prospective error signal in climbing fibers. Program No. 579.01. Neuroscience 2019 Abstracts. Chicago, IL: Society for Neuroscience, 2019. Online.
- Okubo Y, Kakizawa S, Hirose K, Iino M (2004) Cross talk between metabotropic and ionotropic glutamate receptor-mediated signaling in parallel fiber-induced inositol 1,4,5-trisphosphate production in cerebellar Purkinje cells. *J Neurosci* 24:9513–9520. <https://doi.org/10.1523/JNEUROSCI.1829-04.2004>.
- Pasalar S, Roitman AV, Durfee WK, Ebner TJ (2006) Force field effects on cerebellar Purkinje cell discharge with implications for internal models. *Nat Neurosci* 9:1404–1411. <https://doi.org/10.1038/nn1783>.
- Payne HL, French RL, Guo CC, Barbara Nguyen-Vu TD, Manninen T, Raymond JL (2019) Cerebellar Purkinje cells control eye movements with a rapid rate code that is invariant to spike irregularity. *eLife* 8: e37102. doi: 10.7554/eLife.37102.
- Perkel DJ, Hestrin S, Sah P, Nicoll RA (1990) Excitatory synaptic currents in Purkinje cells. *Proc Biol Sci* 241:116–121. <https://doi.org/10.1098/rspb.1990.0074>.
- Ramrani N (2006) The primate cortico-cerebellar system: anatomy and function. *Nat Rev Neurosci* 7(7):511–522. <https://doi.org/10.1038/nrn1953>.
- Rancz EA, Ishikawa T, Duguid I, Chadderton P, Mahon S, Häusser M (2007) High-fidelity transmission of sensory information by single cerebellar mossy fibre boutons. *Nature* 450(7173):1245–1248. <https://doi.org/10.1038/nature05995>.
- Rasmussen A, Jirenhed DA, Hessel G (2008) Simple and complex spike firing patterns in Purkinje cells during classical conditioning. *Cerebellum* 7(4):563–566. <https://doi.org/10.1007/s12311-008-0068-2>.
- Rasmussen A, Jirenhed DA, Zucca R, Johansson F, Svensson P, Hessel G (2013) Number of spikes in climbing fibers determines the direction of cerebellar learning. *J Neurosci* 33(33):13436–13440. <https://doi.org/10.1523/JNEUROSCI.1527-13.2013>.
- Raymond JL, Medina JF (2018) Computational principles of supervised learning in the cerebellum. *Annu Rev Neurosci* 41:233–253. <https://doi.org/10.1146/annurev-neuro-080317-061948>.
- Renzi M, Farrant M, Cull-Candy SG (2007) Climbing-fibre activation of NMDA receptors in Purkinje cells of adult mice. *J Physiol* 585:91–101. <https://doi.org/10.1113/jphysiol.2007.141531>.
- Requarth T, Sawtell NB (2011) Neural mechanisms for filtering self-generated sensory signals in cerebellum-like circuits. *Curr Opin Neurobiol* 21(4):602–608. <https://doi.org/10.1016/j.conb.2011.05.031>.
- Rowan MJM, Bonnan A, Zhang K, Amat SB, Kikuchi C, Taniguchi H, Augustine GJ, Christie JM (2018) Graded control of climbing-fiber-mediated plasticity and learning by inhibition in the cerebellum. *Neuron* 99(5):999–1015.e6. <https://doi.org/10.1016/j.neuron.2018.07.024>.
- Sakurai M (1987) Synaptic modification of parallel fibre-Purkinje cell synapse transmission in in vitro guinea-pig cerebellar slices. *J Physiol* 394:463–480.
- Samejima K, Katagiri K, Doya K, Kawato M (2006) Sybolization and imitation learning of motion sequence using competitive modules. *Electr Commun Japan, Part III* 89(9):42–53. <https://doi.org/10.1002/ecjc.20267>.
- Sanger TD, Kawato M (2020) A cerebellar computational mechanism for delay conditioning at precise time intervals. *Neural Compu submitted*, in press.
- Sanger TD, Yamashita O, Kawato M (2020) Expansion coding and computation in the cerebellum: 50 years after the Marr-Albus codon theory. *J Physiol* 598(5):913–928. <https://doi.org/10.1113/JP278745>.
- Sawtell NB (2017) Neural mechanisms for predicting the sensory consequences of behavior: insights from electrosensory systems. *Annu Rev Physiol* 79:381–399. <https://doi.org/10.1146/annurev-physiol-021115-105003>.
- Schmahmann JD, Guell X, Stoodley CJ, Halko MA (2019) The theory and neuroscience of cerebellar cognition. *Annu Rev Neurosci* 42:337–364. <https://doi.org/10.1146/annurev-neuro-070918-050258>.
- Schmidt-Hieber J (2017) Nonparametric regression using deep neural networks with ReLU activation function. arXiv:1708.06633.
- Schonewille M, Gao Z, Boele HJ, Veloz MF, Amerika WE, Simek AA, De Jeu MT, Steinberg JP, et al. (2011) Reevaluating the role of LTD in cerebellar motor learning. *Neuron* 70(1):43–50. <https://doi.org/10.1016/j.neuron.2011.02.044>.

- Schweighofer N, Doya K, Lay F (2001) Unsupervised learning of granule cell sparse codes enhances cerebellar adaptive control. *Neuroscience* 103(1):35–50. [https://doi.org/10.1016/s0306-4522\(00\)00548-0](https://doi.org/10.1016/s0306-4522(00)00548-0).
- Sendhilnathan N, Ipata AE, Goldberg ME (2020) Neural correlates of reinforcement learning in mid-lateral cerebellum. *Neuron* (19):31098. <https://doi.org/10.1016/j.neuron.2019.12.032>. pii: S0896-6273(19)31098-0.
- Sgritta M, Locatelli F, Soda T, Prestori F, D'Angelo EU (2017) Hebbian spike-timing dependent plasticity at the cerebellar input stage. *J Neurosci* 37(11):2809–2823. <https://doi.org/10.1523/JNEUROSCI.2079-16.2016>.
- Shen W, Flajolet M, Greengard P, Surmeier DJ (2008) Dichotomous dopaminergic control of striatal synaptic plasticity. *Science* 321:848–851. <https://doi.org/10.1126/science.1160575>.
- Shepherd GM, Rowe TB (2017) Neocortical lamination: insights from neuron types and evolutionary precursors. *Front. Neuroanat* 11:100. <https://doi.org/10.3389/fnana.2017.00100>.
- Shidara M, Kawano K, Gomi H, Kawato M (1993) Inverse-dynamics model eye movement control by Purkinje cells in the cerebellum. *Nature* 365(6441):50–52. <https://doi.org/10.1038/365050a0>.
- Shimuta M, Sugihara I, Ishikawa T (2019) Convergence of unisensory-evoked signals via multiple pathways to the cerebellum. *bioRxiv* 10.1101/769000. doi.org/10.1101/769000.
- Shutof F, Ohki M, Kitazawa H, Itohara S, Nagao S (2006) Memory trace of motor learning shifts transsynaptically from cerebellar cortex to nuclei for consolidation. *Neuroscience* 139(2):767–777. <https://doi.org/10.1016/j.neuroscience.2005.12.035>.
- Sillitoe RV, Joyner AL (2007) Morphology, molecular codes, and circuitry produce the three-dimensional complexity of the cerebellum. *Annu Rev Cell Dev Biol* 23:549–577. <https://doi.org/10.1146/annurev.cellbio.23.090506.123237>.
- Simpson JI, Alley KE (1974) Visual climbing fiber input to rabbit vestibulo-cerebellum: a source of direction-specific information. *Brain Res* 82(2):302–308. [https://doi.org/10.1016/0006-8993\(74\)90610-6](https://doi.org/10.1016/0006-8993(74)90610-6).
- Simpson JI, Wylie D, De Zeeuw C (1996) On climbing fiber signals and their consequence (s). *Behav Brain Sci* 19(3):384–398.
- Sokolov AA, Miall RC, Ivry RB (2017) The cerebellum: adaptive prediction for movement and cognition. *Trends Cogn Sci* 21(5):313–332. <https://doi.org/10.1016/j.tics.2017.02.005>.
- Spanne A, Jörntell H (2015) Questioning the role of sparse coding in the brain. *Trends Neurosci* 38(7):417–427. <https://doi.org/10.1016/j.tins.2015.05.005>.
- Streng ML, Popa LS, Ebner TJ (2018) Modulation of sensory prediction error in Purkinje cells during visual feedback manipulations. *Nat Commun* 9(1):1099. <https://doi.org/10.1038/s41467-018-03541-0>.
- Strick PL, Dum RP, Fiez JA (2009) Cerebellum and nonmotor function. *Annu Rev Neurosci*:413–434. <https://doi.org/10.1146/annurev.neuro.31.060407.125606>.
- Stuart G, Häusser M (1994) Initiation and spread of sodium action potentials in cerebellar Purkinje cells. *Neuron* 13:703–712. [https://doi.org/10.1016/0896-6273\(94\)90037-x](https://doi.org/10.1016/0896-6273(94)90037-x).
- Sugimoto N, Samejima K, Doya K, Kawato M (2006) Hierarchical reinforcement learning: temporal abstraction based on MOSAIC model. *IEICE Trans Inf Systems (Japanese edition)* 89(7):1577–1587.
- Sugimoto N, Haruno M, Doya K, Kawato M (2012a) MOSAIC for multiple-reward environments. *Neural Comput* 24:577–606. https://doi.org/10.1162/NECO_a_00246.
- Sugimoto N, Morimoto J, Hyon S, Kawato M (2012b) The eMOSAIC model for humanoid robot control. *Neural Networks* 29–30:8–19. <https://doi.org/10.1016/j.neunet.2012.01.002>.
- Sultan F, Augath M, Hamodeh S, Murayama Y, Oeltermann A, Rauch A, Thier P (2012) Unravelling cerebellar pathways with high temporal precision targeting motor and extensive sensory and parietal networks. *Nat Commun* 3:924. <https://doi.org/10.1038/ncomms1912>.
- Suvrathan A, Payne HL, Raymond JL (2016) Timing rules for synaptic plasticity matched to behavioral function. *Neuron* 92(5):959–967. <https://doi.org/10.1016/j.neuron.2016.10.022>.
- Suvrathan A, Raymond JL (2018) Depressed by learning-heterogeneity of the plasticity rules at parallel fiber synapses onto Purkinje cells. *Cerebellum* 17(6):747–755. <https://doi.org/10.1007/s12311-018-0968-8>.
- Suzuki T (2018) Adaptivity of deep ReLU network for learning in Besov and mixed smooth Besov spaces: optimal rate and curse of dimensionality. arXiv:1810.08033.
- Tabata H, Yamamoto K, Kawato M (2002) Computational study on monkey VOR adaptation and smooth pursuit based on the parallel control-pathway theory. *J Neurophysiol* 87(4):2176–2189. <https://doi.org/10.1152/jn.00168.2001>.
- Takemura A, Ofuji T, Miura K, Kawano K (2017) Neural activity in the dorsal medial superior temporal area of monkeys represents retinal error during adaptive motor learning. *Sci Rep* 7:40939. <https://doi.org/10.1038/srep40939>.
- Tanaka K, Khiroug L, Santamaria F, Doi T, Ogasawara H, Ellis-Davies GC, Kawato M, Augustine GJ (2007) Ca²⁺ requirements for cerebellar long-term synaptic depression: role for a postsynaptic leaky integrator. *Neuron* 54(5):787–800. <https://doi.org/10.1016/j.neuron.2007.05.014>.
- Tang T, Xiao J, Suh CY, Burroughs A, Cerminara NL, Jia L, Marshall SP, Wise AK, et al. (2017) Heterogeneity of Purkinje cell simple spike-complex spike interactions: zebrin- and non-zebrin-related variations. *J Physiol* 595(15):5341–5357. <https://doi.org/10.1113/JP274252>.
- Todorov E, Jordan M (2002) Optimal feedback control as a theory of motor coordination. *Nat Neurosci* 11:1226–1235. <https://doi.org/10.1038/nn963>.
- Tokuda IT, Han CE, Aihara K, Kawato M, Schweighofer N (2010) The role of chaotic resonance in cerebellar learning. *Neural Netw* 23(7):836–842. <https://doi.org/10.1016/j.neunet.2010.04.006>.
- Tokuda IT, Hoang H, Schweighofer N, Kawato M (2013) Adaptive coupling of inferior olive neurons in cerebellar learning. *Neural Netw* 47:42–50. <https://doi.org/10.1016/j.neunet.2012.12.006>.
- Tokuda IT, Hoang H, Kawato M (2017) New insights into olivocerebellar circuits for learning from a small training sample. *Curr Opin Neurobiol* 46:58–67. <https://doi.org/10.1016/j.conb.2017.07.010>.
- Tosches MA, Yamawaki TM, Naumann RK, Jacobi AA, Tushev G, Laurent G (2018) Evolution of pallium, hippocampus, and cortical cell types revealed by single-cell transcriptomics in reptiles. *Science* 360(6391):881–888. <https://doi.org/10.1126/science.aar4237>.
- Tseng Y, Diedrichsen J, Krakauer JW, Shadmehr R, Bastian AJ (2007) Sensory prediction errors drive cerebellum-dependent adaptation of reaching. *J Neurophysiol* 98(1):54–62. <https://doi.org/10.1152/jn.00266.2007>.
- Tsutsumi S, Hidaka N, Isomura Y, Matsuzaki M, Sakimura K, Kano M, Kitamura K (2019) Modular organization of cerebellar climbing fiber inputs during goal-directed behavior. *eLife* 8:e47021. doi: 10.7554/eLife.47021.
- Tsutsumi S, Yamazaki M, Miyazaki T, Watanabe M, Sakimura K, Kano M, Kitamura K (2015) Structure–function relationships between aldolase C/zebrin II expression and complex spike synchrony in the cerebellum. *J Neurosci* 35(2):843–852. <https://doi.org/10.1523/JNEUROSCI.2170-14.2015>.
- Tyrrell T, Willshaw D (1992) Cerebellar cortex: its simulation and the relevance of Marr's theory. *Phil Trans R Soc Lond B* 336(1277):239–257. <https://doi.org/10.1098/rstb.1992.0059>.
- Urakubo H, Honda M, Tanaka K, Kuroda S (2009) Experimental and computational aspects of signaling mechanisms of spike-timing-dependent plasticity. *HFSP J* 3:240–254. <https://doi.org/10.2976/1.3137602>.
- Van Beugen BJ, Gao Z, Boele H-J, Hoebeek F, De Zeeuw CI (2013) High frequency burst firing of granule cells ensures transmission at the parallel fiber to Purkinje cell synapse at the cost of temporal coding. *Front Neural Circuit* 7:95. <https://doi.org/10.3389/fncir.2013.00095>.
- Vetter P, Roth A, Häusser M (2001) Propagation of action potentials in dendrites depends on dendritic morphology. *J Neurophysiol* 85:926–937. <https://doi.org/10.1152/jn.2001.85.2.926>.

- Wagner MJ, Kim TH, Kadmon J, Nguyen ND, Ganguli S, Schnitzer MJ, Luo L (2019) Shared cortex-cerebellum dynamics in the execution and learning of a motor task. *Cell* 177(3):669–682.e24. <https://doi.org/10.1016/j.cell.2019.02.019>.
- Wagner MJ, Kim TH, Savall J, Schnitzer MJ, Luo L (2017) Cerebellar granule cells encode the expectation of reward. *Nature* 544(7648):96–100. <https://doi.org/10.1038/nature21726>.
- Wang SS, Denk W, Häusser M (2000) Coincidence detection in single dendritic spines mediated by calcium release. *Nat Neurosci* 3:1266–1273. <https://doi.org/10.1038/81792>.
- Watanabe S (2009) *Algebraic geometry and statistical learning theory* (Cambridge monographs on applied and computational mathematics). Cambridge University Press. doi: 10.1017/CBO9780511800474.
- Welsh JP, Yamaguchi H, Zeng XH, Kojo M, Nakada Y, Takagi A, Sugimori M, Llinás RR (2005) Normal motor learning during pharmacological prevention of Purkinje cell long-term depression. *Proc Natl Acad Sci U S A* 102(47):17166–17171. <https://doi.org/10.1073/pnas.0508191102>.
- Wickens JR, Begg AJ, Arbutnott GW (1996) Dopamine reverses the depression of rat corticostriatal synapses which normally follows high-frequency stimulation of cortex in vitro. *Neuroscience* 70:1–5. [https://doi.org/10.1016/0306-4522\(95\)00436-m](https://doi.org/10.1016/0306-4522(95)00436-m).
- Widrow B, McCool JM, Larimore MG, Johnson CR (1976) Stationary and nonstationary learning characteristics of the LMS adaptive filter. In: Proceedings of the IEEE, vol. 64, issue. 8, pp 1151–1162. IEEE Institute of electrical and electronics engineering, inc.
- Wolpert DM, Doya K, Kawato M (2003) A unifying computational framework for motor control and social interaction. *Philos Trans Royal Soc: Biol Sci* 358:593–602. <https://doi.org/10.1098/rstb.2002.1238>.
- Wolpert DM, Ghahramani Z, Jordan MI (1995) An internal model for sensorimotor integration. *Science* 269:1880–1882. <https://doi.org/10.1126/science.7569931>.
- Wolpert DM, Kawato M (1998) Multiple paired forward and inverse models for motor control. *Neural Networks* 11(7–8):1317–1329. [https://doi.org/10.1016/s0893-6080\(98\)00066-5](https://doi.org/10.1016/s0893-6080(98)00066-5).
- Wolpert DM, Miall C, Kawato M (1998) Internal models in the cerebellum. *Trends Cognitive Sci* 2:338–347. [https://doi.org/10.1016/s1364-6613\(98\)01221-2](https://doi.org/10.1016/s1364-6613(98)01221-2).
- Xiao J, Cerminara NL, Kotsurovskyy Y, Aoki H, Burroughs A, Wise AK, Luo Y, Marshall SP, et al. (2014) Systematic regional variations in Purkinje cell spiking patterns. *PLoS One* 9(8). doi: 10.1371/journal.pone.0105633. eCollection 2014 e105633.
- Yamada T, Yang Y, Valnegri P, Juric I, Abnoui A, Markwalter KH, Guthrie AN, Godec A, Oldenborg A, Hu M, Holy TE, Bonni A (2019) Sensory experience remodels genome architecture in neural circuit to drive motor learning. *Nature* 569(7758):708–713. <https://doi.org/10.1038/s41586-019-1190-7>.
- Yamaguchi K, Itohara S, Ito M (2016) Reassessment of long-term depression in cerebellar Purkinje cells in mice carrying mutated GluA2 C terminus. *Proc Natl Acad Sci U S A* 113(36):10192–10197. <https://doi.org/10.1073/pnas.1609957113>.
- Yamamoto K, Kawato M, Kotoaska S, Kitazawa S (2007) Encoding of movement dynamics by Purkinje cell simple spike activity during fast arm movements under resistive and assistive force fields. *J Neurophysiol* 97:1588–1599. <https://doi.org/10.1152/jn.00206.2006>.
- Yamamoto K, Kobayashi Y, Takemura A, Kawano K, Kawato M (2002) Computational studies on acquisition and adaptation of ocular following responses based on cerebellar synaptic plasticity. *J Neurophysiol* 87:1554–1571. <https://doi.org/10.1152/jn.00166.2001>.
- Yamazaki T, Nagao S, Lennon W, Tanaka S (2015) Modeling memory consolidation during posttraining periods in cerebellovestibular learning. *Proc Natl Acad Sci U S A* 112(11):3541–3546. <https://doi.org/10.1073/pnas.1413798112>.
- Yamazaki T, Lennon W (2019) Revisiting a theory of cerebellar cortex. *Neurosci Res* 148:1–8. <https://doi.org/10.1016/j.neures.2019.03.001>.
- Yang Y, Lisberger SG (2014) Purkinje cell plasticity and cerebellar motor learning are graded by complex-spike duration. *Nature* 510(7506):529–532. <https://doi.org/10.1038/nature13282>.
- Yang Y, Yamada T, Hill KK, Hemberg M, Reddy NC, Cho HY, Guthrie AN, Oldenborg A, et al. (2016) Chromatin remodeling inactivates activity genes and regulates neural coding. *Science* 353(6296):300–305. <https://doi.org/10.1126/science.aad4225>.
- Zhang C, Bengio S, Hardt M, Recht B, Vinyals O (2017) Understanding deep learning requires rethinking generalization. arXiv:1611.03530.
- Zhang W, Linden DJ (2006) Long-term depression at the mossy fiber–deep cerebellar nucleus synapse. *J Neurosci* 26(26):6935–6944. <https://doi.org/10.1523/JNEUROSCI.0784-06.2006>.
- Zhou H, Lin Z, Voges K, Ju C, Gao Z, Bosman LW, Ruigrok TJ, Hoebeek FE, De Zeeuw CI, Schonewille M (2014) Cerebellar modules operate at different frequencies. *eLife* 3:e02536. doi: 10.7554/eLife.02536.

(Received 6 March 2020, Accepted 15 June 2020)
(Available online 27 June 2020)